

Mouse anti-Human Cytokeratin 8 Monoclonal Antibody | anti-KRT8 antibody

Anti-Cytokeratin 8 Antibody

CATALOG # AAA264953	HOST Mouse	REACTIVITY Human	APPLICATIONS Immunohistochemistry, Western Blot
-------------------------------	----------------------	----------------------------	---

PRODUCT SYNONYMS

Cytokeratin 8; Monoclonal Antibody; Anti-Cytokeratin 8 Antibody; CYK8; Keratin type II cytoskeletal 8; Cytokeratin-8; CK-8; Keratin-8; K8; Type-II keratin Kb8; anti-KRT8 antibody

PRODUCT SPECIFICATIONS


Host	Mouse
Reactivity	Human
Clonality	Monoclonal
Isotype	Mouse IgG2a
Specificity	Recognizes endogenous levels of Cytokeratin 8 protein.
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in PBS containing 50% glycerol, 0.2% BSA and 0.01% sodium azide.
Applicable Applications for anti-KRT8 antibody	IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within human Cytokeratin 8. The exact sequence is proprietary.
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C for one year. Avoid freeze/thaw cycles.

IMAGES




IP (Immunoprecipitation)

Immunohistochemical analysis of Cytokeratin 8 staining in human kidney formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



IHC (Immunohistochemistry)

Immunohistochemical analysis of Cytokeratin 8 staining in human gastric adenocarcinoma formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



IF (Immunofluorescence)

Immunohistochemical analysis of Cytokeratin 8 staining in human hepatocellular carcinoma formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

ADDITIONAL INFORMATION

Related Product Information for anti-KRT8 antibody	Mouse monoclonal antibody to Cytokeratin 8
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	372466577
NCBI GeneID	3856
NCBI Accession #	NP_001243222.1
NCBI GenBank Nucleotide #	NM_001256293.1
UniProt Accession #	P05787; Q14099; Q14716; Q14717; Q53GJ0; Q6DHW5; Q6GMY0; Q6P4C7; Q96J60; A8K4H3; B0AZN5; F8VXB4
Molecular Weight	53,704 Da
NCBI Official Full Name	keratin, type II cytoskeletal 8 isoform 2
NCBI Official Synonym Full Names	keratin 8
NCBI Official Symbol	KRT8
NCBI Official Synonym Symbols	K8; KO; CK8; CK-8; CYK8; K2C8; CARD2
NCBI Protein Information	keratin, type II cytoskeletal 8; cytokeratin 8; cytokeratin-8; type-II keratin Kb8
UniProt Protein Name	Keratin, type II cytoskeletal 8
UniProt Gene Name	KRT8
UniProt Synonym Gene Names	CYK8; CK-8; K8
UniProt Entry Name	K2C8_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/monoclonal/cytokeratin-8-antibody-immunohistochemistry-western-blot-elisa/264953>