

Rabbit anti-Human MCH Receptor 1 Polyclonal Antibody | anti-MCH antibody

Anti-MCH Receptor 1 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA265292	Rabbit	Human	Flow Cytometry, Functional Assay, Immunohistochemistry, Western Blot

PRODUCT SYNONYMS

MCH Receptor 1; Polyclonal Antibody; Anti-MCH Receptor 1 Antibody; anti-MCH antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Specificity	Recognizes endogenous levels of MCH Receptor 1 protein.
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH7.3, 30% glycerol, and 0.01% sodium azide.
Applicable Applications for anti-MCH antibody	FCM/FACS (Flow Cytometry), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the C-terminal region of human MCH Receptor 1. The exact sequence is proprietary.
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C for one year. Avoid freeze/thaw cycles.

IMAGES




FCM/FACS (Flow Cytometry)

Flow cytometric analysis of K562 cells using Anti-MCH Receptor 1 Antibody. The cells were fixed with 2% paraformaldehyde (10 min) and then permeabilized with 90% methanol for 10 min. The cells were incubated in 2% bovine serum albumin to block non-specific protein-protein interactions followed by the antibody at 37 °C for 60 min. The secondary antibody Goat Anti-Rabbit IgG (H&L) - AF488 was incubated at 37 °C for 40 min. Isotype control antibody (blue line) was used under the same condition.



IHC (Immunohistochemistry)

Immunohistochemical analysis of MCH Receptor 1 staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of MCH Receptor 1 expression in mouse brain (A), rat brain (B) whole cell lysates. (Predicted band size: 45 kD; Observed band size: 50 kD)

ADDITIONAL INFORMATION

Related Product Information for anti-MCH antibody	Rabbit polyclonal antibody to MCH Receptor 1
---	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	88758590
NCBI GeneID	2847
NCBI Accession #	NP_005288.3
NCBI GenBank Nucleotide #	NM_005297.3
UniProt Accession #	Q99705; Q5R3J1; Q96S47; Q9BV08; B2RBX6
Molecular Weight	45,963 Da
NCBI Official Full Name	melanin-concentrating hormone receptor 1
NCBI Official Synonym Full Names	melanin-concentrating hormone receptor 1
NCBI Official Symbol	MCHR1
NCBI Official Synonym Symbols	SLC1; GPR24; MCH1R; SLC-1; MCH-1R
NCBI Protein Information	melanin-concentrating hormone receptor 1; MCH receptor 1; G protein-coupled receptor 24; somatostatin receptor-like protein
UniProt Protein Name	Melanin-concentrating hormone receptor 1
UniProt Gene Name	MCHR1
UniProt Synonym Gene Names	GPR24; SLC1; MCH receptor 1; MCH-R1; MCHR-1; MCH1R; MCHR
UniProt Entry Name	MCHR1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.