Rabbit anti-Human DCAF8L2 Polyclonal Antibody | anti-DCAF8L2 antibody

Anti-DCAF8L2 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA265407	**Rabbit**	**Human**	**Immunofluorescence, Immunohistochemistry, Western Blot**


PRODUCT SYNONYMS

DCAF8L2; Polyclonal Antibody; Anti-DCAF8L2 Antibody; anti-DCAF8L2 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Specificity	Recognizes endogenous levels of DCAF8L2 protein.
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH7.3, 30% glycerol, and 0.01% sodium azide.
Applicable Applications for anti-DCAF8L2 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the N-terminal region of human DCAF8L2. The exact sequence is proprietary.
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C for one year. Avoid freeze/thaw cycles.

IMAGES



IF (Immunofluorescence)
Immunofluorescent analysis of Anti-DCAF8L2 staining in ZR751 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a hidified chamber. Cells were washed with PBST and incubated with a AF488-conjugated secondary antibody (green) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).



IHC (Immunohiostchemistry)
Immunohistochemical analysis of DCAF8L2 staining in human breast carcinoma formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)
Western blot analysis of DCAF8L2 expression in K562 (A) whole cell lysates. (Predicted band size: 71 kD; Observed band size: 71 kD

ADDITIONAL INFORMATION

Related Product Information for anti-DCAF8L2 antibody	Rabbit polyclonal antibody to DCAF8L2

Product Categories/Family for anti-DCAF8L2 antibody	Primary Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	187957316
NCBI GeneID	347442
UniProt Accession #	P0C7V8; B2RXH9; J3KT06
Molecular Weight	71,191 Da
NCBI Official Full Name	WDR42C protein
NCBI Official Synonym Full Names	DDB1 and CUL4 associated factor 8-like 2
NCBI Official Symbol	DCAF8L2
NCBI Official Synonym Symbols	WDR42C
NCBI Protein Information	DDB1- and CUL4-associated factor 8-like protein 2; WD repeat domain 42C; WD repeat-containing protein 42C
UniProt Protein Name	DDB1- and CUL4-associated factor 8-like protein 2
UniProt Gene Name	**DCAF8L2**
UniProt Synonym Gene Names	WDR42C
UniProt Entry Name	DC8L2_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: https://stage.aaabiotech.com/polyclonal/dcaf8l2-antibody-immunofluorescence-immunohistochemistry-western-blot-elisa/265407