

Rabbit AKT Monoclonal Antibody | anti-AKT antibody

Anti-AKT Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA265592	Rabbit	Human, Mouse, Rat, Hamster	Immunoprecipitation, Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

AKT; Monoclonal Antibody; Anti-AKT Antibody; anti-AKT antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat, Hamster
Clonality	Monoclonal
Specificity	Recognizes endogenous levels of AKT protein.
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 50mM Tris-Glycine (pH7.4), 0.15M NaCl, 50% Glycerol, 0.01% Sodium azide and 0.05% BSA.
Applicable Applications for anti-AKT antibody	IP (Immunoprecipitation), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	Recombinant protein of human AKT. The exact sequence is proprietary.
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C for one year. Avoid freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of AKT staining in A549 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a AF488-conjugated secondary antibody (green) in PBS at room temperature in the dark.



IHC (Immunohistochemistry)

Immunohistochemical analysis of AKT staining in human lung cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of AKT expression in Hela (A), Jurkat (B), rat brain (C), CHO-K1 (D) whole cell lysates. (Predicted band size: 55 kD; Observed band size: 55 kD)

ADDITIONAL INFORMATION

Related Product Information for anti-AKT antibody	Rabbit monoclonal antibody to AKT
---	-----------------------------------

NCBI AND UNIPROT INFORMATION

NCBI GI #	62241013
NCBI GeneID	207
NCBI Accession #	NP_001014431.1
NCBI GenBank Nucleotide #	NM_001014431.1
UniProt Accession #	P31749; P31750; Q60823; Q9WUA6; P47196; P47197; Q63484; P31751; Q9Y243
Molecular Weight	55,686 Da
NCBI Official Full Name	RAC-alpha serine/threonine-protein kinase
NCBI Official Synonym Full Names	v-akt murine thymoma viral oncogene homolog 1
NCBI Official Symbol	AKT1
NCBI Official Synonym Symbols	AKT; PKB; RAC; CWS6; PRKBA; PKB-ALPHA; RAC-ALPHA
NCBI Protein Information	RAC-alpha serine/threonine-protein kinase; PKB alpha; RAC-PK-alpha; proto-oncogene c-Akt; protein kinase B alpha; rac protein kinase alpha
UniProt Protein Name	RAC-alpha serine/threonine-protein kinase
UniProt Gene Name	AKT1
UniProt Synonym Gene Names	PKB; RAC; PKB; PKB alpha
UniProt Entry Name	AKT1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/monoclonal/akt-antibody-immunoprecipitation-immunofluorescence-immunohistochemistry-western-blot-elisa/265592>