

Rabbit ERK2 Monoclonal Antibody | anti-ERK2 antibody

Anti-ERK2 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA265619	Rabbit	Human, Mouse, Rat	Immunoprecipitation, Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

ERK2; Monoclonal Antibody; Anti-ERK2 Antibody; anti-ERK2 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Monoclonal
Specificity	Recognizes endogenous levels of ERK2 protein.
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 50mM Tris-Glycine (pH7.4), 0.15M NaCl, 50% Glycerol, 0.01% Sodium azide and 0.05% BSA.
Applicable Applications for anti-ERK2 antibody	IP (Immunoprecipitation), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	A synthetic peptide of human ERK2
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C for one year. Avoid freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of ERK2 staining in K562 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a AF488-conjugated secondary antibody (green) in PBS at room temperature in the dark.



IHC (Immunohistochemistry)

Immunohistochemical analysis of ERK2 staining in human cholangiocarcinoma formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.41). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of ERK2 expression in K562 (A), rat brain (B), C6 (C), NIH3T3 (D), HeLa (E) whole cell lysates. (Predicted band size: 41 kD; Observed band size: 41 kD)

ADDITIONAL INFORMATION

Related Product Information for anti-ERK2 antibody	Rabbit monoclonal antibody to ERK2
Product Categories/Family for anti-ERK2 antibody	Primary Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	66932916
NCBI GeneID	5594
NCBI Accession #	NP_002736.3
NCBI GenBank Nucleotide #	NM_002745.4
UniProt Accession #	P28482 ; P63085 ; P63086
Molecular Weight	41,390 Da
NCBI Official Full Name	mitogen-activated protein kinase 1
NCBI Official Synonym Full Names	mitogen-activated protein kinase 1
NCBI Official Symbol	MAPK1
NCBI Official Synonym Symbols	ERK; p38; p40; p41; ERK2; ERT1; MAPK2; PRKM1; PRKM2; P42MAPK; p41mapk
NCBI Protein Information	mitogen-activated protein kinase 1; ERK-2; MAPK 2; p42-MAPK; MAP kinase 1; MAP kinase 2; MAP kinase isoform p42; protein tyrosine kinase ERK2; mitogen-activated protein kinase 2; extracellular signal-regulated kinase 2
UniProt Protein Name	Mitogen-activated protein kinase 1
UniProt Gene Name	MAPK1
UniProt Synonym Gene Names	ERK2; PRKM1; PRKM2; MAP kinase 1; MAPK 1; ERK-2; p42-MAPK; MAP kinase 2; MAPK 2
UniProt Entry Name	MK01_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/monoclonal/erk2-antibody-immunoprecipitation-immunofluorescence-immunohistochemistry-western-blot-elisa/265619>