

Rabbit UBC9 Monoclonal Antibody | anti-UBC9 antibody

Anti-UBC9 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA265738	Rabbit	Human, Mouse, Rat	Immunoprecipitation, Immunofluorescence, Immunohistochemistry, Western Blot

PRODUCT SYNONYMS

UBC9; Monoclonal Antibody; Anti-UBC9 Antibody; anti-UBC9 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Monoclonal
Specificity	Recognizes endogenous levels of UBC9 protein.
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 50mM Tris-Glycine (pH7.4), 0.15M NaCl, 50% Glycerol, 0.01% Sodium azide and 0.05% BSA.
Applicable Applications for anti-UBC9 antibody	IP (Immunoprecipitation), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	A synthetic peptide of human UBE2I/UBC9
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C for one year. Avoid freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of UBC9 staining in HeLa cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a AF488-conjugated secondary antibody (green) in PBS at room temperature in the dark.



IHC (Immunohistochemistry)

Immunohistochemical analysis of UBC9 staining in human tonsil formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.132). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of UBC9 expression in Hela (A), A549 (B), HL60 (C), NIH3T3 (D), PC12 (E), MEF (F) whole cell lysates. (Predicted band size: 18 kD; Observed band size: 18 kD)

ADDITIONAL INFORMATION

Related Product Information for anti-UBC9 antibody	Rabbit monoclonal antibody to UBC9
--	------------------------------------

NCBI AND UNIPROT INFORMATION

NCBI GI #	4507785
NCBI GeneID	7329
NCBI Accession #	NP_003336.1
NCBI GenBank Nucleotide #	NM_003345.4
UniProt Accession #	P63279; P63280; P63281
Molecular Weight	158
NCBI Official Full Name	SUMO-conjugating enzyme UBC9
NCBI Official Synonym Full Names	ubiquitin-conjugating enzyme E2I
NCBI Official Symbol	UBE2I
NCBI Official Synonym Symbols	P18; UBC9; C358B7.1
NCBI Protein Information	SUMO-conjugating enzyme UBC9; SUMO-protein ligase; SUMO-1-protein ligase; ubiquitin-protein ligase I; ubiquitin carrier protein 9; ubiquitin carrier protein I; ubiquitin-protein ligase E2I; ubiquitin conjugating enzyme 9; ubiquitin-conjugating enzyme UbcE
UniProt Protein Name	SUMO-conjugating enzyme UBC9
UniProt Gene Name	UBE2I
UniProt Synonym Gene Names	UBC9; UBCE9
UniProt Entry Name	UBC9_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/monoclonal/ubc9-antibody-immunoprecipitation-immunofluorescence-immunohistochemistry-western-blot-elisa/265738>