

Mouse anti-Human Vimentin Monoclonal Antibody | anti-VIM antibody

Anti-Vimentin Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA265800	Mouse	Human	Immunoprecipitation, Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

Vimentin; Monoclonal Antibody; Anti-Vimentin Antibody; anti-VIM antibody

PRODUCT SPECIFICATIONS


Host	Mouse
Reactivity	Human
Clonality	Monoclonal
Specificity	Recognizes endogenous levels of Vimentin protein.
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide, pH7.3.
Applicable Applications for anti-VIM antibody	IP (Immunoprecipitation), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	Purified recombinant human Vimentin protein fragments expressed in E coli.
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C for one year. Avoid freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of Vimentin staining in HeLa cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a AF488-conjugated secondary antibody (green) in PBS at room temperature in the dark.



IHC (Immunohistochemistry)

Immunohistochemical analysis of Vimentin staining in human tonsil formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.136). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of Vimentin expression in Molt4 (A), K562 (B), COS7 (C), Jurkat (D), Hela (E) whole cell lysates. (Predicted band size: 54 kD; Observed band size: 57 kD)

ADDITIONAL INFORMATION

Related Product Information for anti-VIM antibody	Mouse monoclonal antibody to Vimentin
---	---------------------------------------

NCBI AND UNIPROT INFORMATION

NCBI GI #	62414289
NCBI GeneID	7431
NCBI Accession #	NP_003371.2
NCBI GenBank Nucleotide #	NM_003380.3
UniProt Accession #	P08670; Q15867; Q15868; Q15869; Q548L2; Q6LER9; Q8N850; Q96ML2; Q9NTM3; B0YJC2; D3DRU4
Molecular Weight	53,652 Da
NCBI Official Full Name	vimentin
NCBI Official Synonym Full Names	vimentin
NCBI Official Symbol	VIM
NCBI Protein Information	vimentin
UniProt Protein Name	Vimentin
UniProt Gene Name	VIM
UniProt Entry Name	VIME_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/monoclonal/vimentin-antibody-immunoprecipitation-immunofluorescence-immunohistochemistry-western-blot-elisa/265800>