

Mouse anti-Human LCK Monoclonal Antibody | anti-LCK antibody

Anti-LCK Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA265944	Mouse	Human	Flow Cytometry, Functional Assay, Immunofluorescence, Immunohistochemistry, Western Blot

PRODUCT SYNONYMS

LCK; Monoclonal Antibody; Anti-LCK Antibody; anti-LCK antibody

PRODUCT SPECIFICATIONS

Host	Mouse
Reactivity	Human
Clonality	Monoclonal
Isotype	Mouse IgG1 kappa
Specificity	Recognizes endogenous levels of LCK protein.
Purity/Purification	This antibody is purified through a protein G column.
Form/Format	Mouse IgG1 kappa. Liquid in PBS, pH7.3, 30% glycerol, and 0.01% sodium azide.
Applicable Applications for anti-LCK antibody	FCM/FACS (Flow Cytometry), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	Recombinant fusion protein of human LCK. The exact sequence is proprietary.
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C for one year. Avoid freeze/thaw cycles.

IMAGES




FCM/FACS (Flow Cytometry)

Flow cytometric analysis of Jurkat cells using Anti-LCK Antibody. The cells were fixed with 2% paraformaldehyde (10 min) and then permeabilized with 90% methanol for 10 min. The cells were incubated in 2% bovine serum albumin to block non-specific protein-protein interactions followed by the antibody at 37 °C for 60 min. The secondary antibody Goat Anti-Mouse IgG (H&L) - AF488 was incubated at 37 °C for 40 min. Isotype control antibody (blue line) was used under the same condition.




IF (Immunofluorescence)

Immunofluorescent analysis of LCK staining in Hela cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a AF488-conjugated secondary antibody (green) in PBS at room temperature in the dark. Phalloidin - AF555 was used to stain Actin filaments (red). DAPI was used to stain the cell nuclei (blue).



IHC (Immunohistochemistry)

Immunohistochemical analysis of LCK staining in human thymus formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of LCK expression in Jurkat (A) whole cell lysates. (Predicted band size: 58; 41 kD; Observed band size: 41 kD)

ADDITIONAL INFORMATION

Related Product Information for anti-LCK antibody	Mouse monoclonal antibody to LCK
Product Categories/Family for anti-LCK antibody	Primary Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	112789548
NCBI GeneID	3932
NCBI Accession #	NP_001036236.1
NCBI GenBank Nucleotide #	NM_001042771.1
UniProt Accession #	P06239; P07100; Q12850; Q13152; Q5TDH8; Q5TDH9; Q7RTZ3; Q96DW4; Q9NYT8; D3DPP8
Molecular Weight	58,001 Da
NCBI Official Full Name	tyrosine-protein kinase Lck
NCBI Official Synonym Full Names	lymphocyte-specific protein tyrosine kinase
NCBI Official Symbol	LCK
NCBI Official Synonym Symbols	LSK; YT16; p56lck; pp58lck
NCBI Protein Information	tyrosine-protein kinase Lck; leukocyte C-terminal Src kinase; p56(LSTRA) protein-tyrosine kinase; t cell-specific protein-tyrosine kinase; proto-oncogene tyrosine-protein kinase LCK; lymphocyte cell-specific protein-tyrosine kinase; T-lymphocyte specific
UniProt Protein Name	Tyrosine-protein kinase Lck
UniProt Gene Name	LCK
UniProt Synonym Gene Names	LSK
UniProt Entry Name	LCK_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/monoclonal/lck-antibody-flow-cytometry-immunofluorescence-immunohistochemistry-western-blot-elisa/265944>