

Mouse anti-Human, Mouse CD271 Monoclonal Antibody | anti-CD271 antibody

Anti-CD271 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA265957	Mouse	Human, Mouse	Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

CD271; Monoclonal Antibody; Anti-CD271 Antibody; anti-CD271 antibody

PRODUCT SPECIFICATIONS


Host	Mouse
Reactivity	Human, Mouse
Clonality	Monoclonal
Isotype	Mouse IgG2b kappa
Specificity	Recognizes endogenous levels of CD271 protein.
Purity/Purification	This antibody is purified through a protein G column.
Form/Format	Mouse IgG2b kappa. Liquid in PBS, pH7.3, 30% glycerol, and 0.01% sodium azide.
Applicable Applications for anti-CD271 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	Recombinant fusion protein of human CD271. The exact sequence is proprietary.
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C for one year. Avoid freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of CD271 staining in SHSY5Y cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a AF488-conjugated secondary antibody (green) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).



IHC (Immunohistochemistry)

Immunohistochemical analysis of CD271 staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of CD271 expression in mouse heart (A) whole cell lysates. (Predicted band size: 45 kD; Observed band size: 55 kD)

ADDITIONAL INFORMATION

Related Product Information for anti-CD271 antibody	Mouse monoclonal antibody to CD271
Product Categories/Family for anti-CD271 antibody	Primary Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	4505393
NCBI GeneID	4804
NCBI Accession #	NP_002498.1
NCBI GenBank Nucleotide #	NM_002507.3
UniProt Accession #	P08138 ; Q9Z0W1
Molecular Weight	45,183 Da
NCBI Official Full Name	tumor necrosis factor receptor superfamily member 16
NCBI Official Synonym Full Names	nerve growth factor receptor
NCBI Official Symbol	NGFR
NCBI Official Synonym Symbols	CD271; p75NTR; TNFRSF16; p75(NTR); Gp80-LNGFR
NCBI Protein Information	tumor necrosis factor receptor superfamily member 16; p75 ICD; NGF receptor; TNFR superfamily, member 16; low affinity nerve growth factor receptor; low affinity neurotrophin receptor p75NTR; low-affinity nerve growth factor receptor
UniProt Protein Name	Tumor necrosis factor receptor superfamily member 16
UniProt Gene Name	NGFR
UniProt Synonym Gene Names	TNFRSF16; NGF receptor
UniProt Entry Name	TNR16_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/monoclonal/cd271-antibody-immunofluorescence-immunohistochemistry-western-blot-elisa/265957>