

Mouse anti-Human CD44 Monoclonal Antibody | anti-CD44 antibody

Anti-CD44 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA265988	Mouse	Human	Flow Cytometry, Functional Assay, Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

CD44; Monoclonal Antibody; Anti-CD44 Antibody; anti-CD44 antibody

PRODUCT SPECIFICATIONS


Host	Mouse
Reactivity	Human
Clonality	Monoclonal
Isotype	Mouse IgG2a kappa
Specificity	Recognizes endogenous levels of CD44 protein.
Purity/Purification	This antibody is purified through a protein G column.
Form/Format	Mouse IgG2a kappa. Liquid in PBS, pH7.3, 30% glycerol, and 0.01% sodium azide.
Applicable Applications for anti-CD44 antibody	FCM/FACS (Flow Cytometry), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	Recombinant fusion protein of human CD44. The exact sequence is proprietary.
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C for one year. Avoid freeze/thaw cycles.

IMAGES




FCM/FACS (Flow Cytometry)

Flow cytometric analysis of Hela cells using Anti-CD44 Antibody. The cells were fixed with 2% paraformaldehyde (10 min) and then permeabilized with 90% methanol for 10 min. The cells were incubated in 2% bovine serum albumin to block non-specific protein-protein interactions followed by the antibody at 37 °C for 60 min. The secondary antibody Goat Anti-Mouse IgG (H&L) - AF488 was incubated at 37 °C for 40 min. Isotype control antibody (blue line) was used under the same condition.




IF (Immunofluorescence)

Immunofluorescent analysis of CD44 staining in Hela cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a AF594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).



IHC (Immunohistochemistry)

Immunohistochemical analysis of CD44 staining in human esophagus carcinoma formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of CD44 expression in Hela (A) whole cell lysates. (Predicted band size: 81 kD; Observed band size: 90 kD)

ADDITIONAL INFORMATION

Related Product Information for anti-CD44 antibody	Mouse monoclonal antibody to CD44
Product Categories/Family for anti-CD44 antibody	Primary Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	48255935
NCBI GeneID	960
NCBI Accession #	NP_000601.3
NCBI GenBank Nucleotide #	NM_000610.3
UniProt Accession #	P16070; O95370; P22511; Q04858; Q13419; Q13957; Q13958; Q13959; A5YRN9; B6EAT9; D3DR12; D3DR13
Molecular Weight	81,538 Da
NCBI Official Full Name	CD44 antigen isoform 1
NCBI Official Synonym Full Names	CD44 molecule (Indian blood group)
NCBI Official Symbol	CD44
NCBI Official Synonym Symbols	IN; LHR; MC56; MDU2; MDU3; MIC4; Pgp1; CDW44; CSPG8; HCELL; HUTCH-I; ECMR-III
NCBI Protein Information	CD44 antigen; epican; Hermes antigen; hyaluronate receptor; phagocytic glycoprotein 1; heparan sulfate proteoglycan; cell surface glycoprotein CD44; extracellular matrix receptor III; chondroitin sulfate proteoglycan 8; GP90 lymphocyte homing/adhesion rec
UniProt Protein Name	CD44 antigen
UniProt Gene Name	CD44
UniProt Synonym Gene Names	LHR; MDU2; MDU3; MIC4; ECMR-III; PGP-1; PGP-I
UniProt Entry Name	CD44_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.