

Mouse anti-Human CD49f Monoclonal Antibody | anti-CD49f antibody

Anti-CD49f Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA266005	Mouse	Human	Flow Cytometry, Functional Assay, Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

CD49f; Monoclonal Antibody; Anti-CD49f Antibody; anti-CD49f antibody

PRODUCT SPECIFICATIONS


Host	Mouse
Reactivity	Human
Clonality	Monoclonal
Isotype	Mouse IgG1 kappa
Specificity	Recognizes endogenous levels of CD49f protein.
Purity/Purification	This antibody is purified through a protein G column.
Form/Format	Mouse IgG1 kappa. Liquid in PBS, pH7.3, 30% glycerol, and 0.01% sodium azide.
Applicable Applications for anti-CD49f antibody	FCM/FACS (Flow Cytometry), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	Recombinant fusion protein of human CD49f. The exact sequence is proprietary.
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C for one year. Avoid freeze/thaw cycles.

IMAGES




FCM/FACS (Flow Cytometry)

Flow cytometric analysis of HepG2 cells using Anti-CD49f Antibody. The cells were fixed with 2% paraformaldehyde (10 min) and then permeabilized with 90% methanol for 10 min. The cells were incubated in 2% bovine serum albumin to block non-specific protein-protein interactions followed by the antibody at 37 °C for 60 min. The secondary antibody Goat Anti-Mouse IgG (H&L) - AF488 was incubated at 37 °C for 40 min. Isotype control antibody (blue line) was used under the same condition.




IF (Immunofluorescence)

Immunofluorescent analysis of CD49f staining in HepG2 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a AF488-conjugated secondary antibody (green) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).



IHC (Immunohistochemistry)

Immunohistochemical analysis of CD49f staining in human skin carcinoma formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of CD49f expression in HepG2 (A) whole cell lysates. (Predicted band size: 126 kD; Observed band size: 126 kD)

ADDITIONAL INFORMATION

Related Product Information for anti-CD49f antibody	Mouse monoclonal antibody to CD49f
Product Categories/Family for anti-CD49f antibody	Primary Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	119395742
NCBI GeneID	3655
NCBI Accession #	NP_000201.2
NCBI GenBank Nucleotide #	NM_000210.2
UniProt Accession #	P23229 ; Q08443 ; Q0MRC7 ; Q14646 ; Q16508 ; Q53RX7 ; Q59HB7 ; Q86VL6 ; B2RMU9 ; B4DG69 ; B4DKB8 ; C4AM96
Molecular Weight	126,633 Da
NCBI Official Full Name	integrin alpha-6 isoform b
NCBI Official Synonym Full Names	integrin, alpha 6
NCBI Official Symbol	ITGA6
NCBI Official Synonym Symbols	CD49f; VLA-6; ITGA6B
NCBI Protein Information	integrin alpha-6; integrin alpha6B; CD49 antigen-like family member F
UniProt Protein Name	Integrin alpha-6
UniProt Gene Name	ITGA6
UniProt Entry Name	ITA6_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/monoclonal/cd49f-antibody-flow-cytometry-immunofluorescence-immunohistochemistry-western-blot-elisa/266005>