

Mouse anti-Human, Mouse ERK1 Monoclonal Antibody | anti-ERK1 antibody

Anti-ERK1 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA266011	Mouse	Human, Mouse	Immunofluorescence, Immunohistochemistry, Western Blot

PRODUCT SYNONYMS

ERK1; Monoclonal Antibody; Anti-ERK1 Antibody; anti-ERK1 antibody

PRODUCT SPECIFICATIONS

Host	Mouse
Reactivity	Human, Mouse
Clonality	Monoclonal
Isotype	Mouse IgG1 kappa
Specificity	Recognizes endogenous levels of ERK1 protein.
Purity/Purification	This antibody is purified through a protein G column.
Form/Format	Mouse IgG1 kappa. Liquid in PBS, pH7.3, 30% glycerol, and 0.01% sodium azide.
Applicable Applications for anti-ERK1 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human ERK1. The exact sequence is proprietary.
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C for one year. Avoid freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of ERK1 staining in Hela cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a AF488-conjugated secondary antibody (green) in PBS at room temperature in the dark. Phalloidin - AF555 was used to stain Actin filaments (red). DAPI was used to stain the cell nuclei (blue).



IHC (Immunohistochemistry)

Immunohistochemical analysis of ERK1 staining in human breast carcinoma formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of ERK1 expression in MCF7 (A), Hela (B), Jurkat (C), mouse brain (D) whole cell lysates. (Predicted band size: 43 kD; Observed band size: 44 kD)

ADDITIONAL INFORMATION

Related Product Information for anti-ERK1 antibody	Mouse monoclonal antibody to ERK1
Product Categories/Family for anti-ERK1 antibody	Primary Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	91718897
NCBI GeneID	5595
NCBI Accession #	NP_001035145.1
NCBI GenBank Nucleotide #	NM_001040056.2
UniProt Accession #	P27361 ; Q63844
Molecular Weight	43,136 Da
NCBI Official Full Name	mitogen-activated protein kinase 3 isoform 2
NCBI Official Synonym Full Names	mitogen-activated protein kinase 3
NCBI Official Symbol	MAPK3
NCBI Official Synonym Symbols	ERK1; ERT2; ERK-1; PRKM3; P44ERK1; P44MAPK; HS44KDAP; HUMKER1A; p44-ERK1; p44-MAPK
NCBI Protein Information	mitogen-activated protein kinase 3; MAPK 1; MAP kinase isoform p44; insulin-stimulated MAP2 kinase; extracellular signal-related kinase 1; extracellular signal-regulated kinase 1; microtubule-associated protein 2 kinase
UniProt Protein Name	Mitogen-activated protein kinase 3
UniProt Gene Name	MAPK3
UniProt Synonym Gene Names	ERK1; PRKM3; MAP kinase 3; MAPK 3; ERK-1; p44-MAPK
UniProt Entry Name	MK03_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/monoclonal/erk1-antibody-immunofluorescence-immunohistochemistry-western-blot-elisa/266011>