

IgG secondary antibody

Goat Anti-Mouse/Rabbit IgG (H&L)-HRP Polymer

CATALOG # AAA266236	HOST Goat	REACTIVITY Mouse, Rabbit	APPLICATIONS Immunohistochemistry, Western Blot, ELISA
-------------------------------	---------------------	------------------------------------	--


PRODUCT SYNONYMS

IgG; Polyclonal Secondary Antibody; Goat Anti-Mouse/Rabbit IgG (H&L)-HRP Polymer; IgG secondary antibody

PRODUCT SPECIFICATIONS


Host	Goat
Reactivity	Mouse, Rabbit
Clonality	Polyclonal
Specificity	By immunoelectrophoresis and ELISA this antibody reacts specifically with Mouse IgG and Rabbit IgG. No antibody was detected against non immunoglobulin serum proteins.
Purity/Purification	Goat Polyclonal Secondary Antibody to Mouse/Rabbit IgG (H&L) have been cross-adsorbed against IgG from bovine, goat, horse and human. Cross-adsorption or pre-adsorption is a purification step to increase specificity of the antibody resulting in less backg
Form/Format	Liquid in 0.01M Phosphate Buffered Saline, pH7.2, containing 1% BSA, 50% glycerol, 0.05% Proclin
Concentration	0.5mg/ml (varies by lot)
Applicable Applications for IgG secondary antibody	IHC (Immunohistochemistry), WB (Western Blot), ELISA
Immunogen	Mouse/Rabbit IgG
Conjugation	HRP Polymer
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C for one year. Avoid freeze/thaw cycles.

IMAGES



IHC (Immunohistochemistry)

Immunohistochemical analysis staining in human liver carcinoma formalin fixed paraffin-embedded tissue section. The section was pre-treated using pressure cooker heat antigen retrieval with sodium citrate buffer (0.01M, pH=6) for 3 minutes. The section was detected using rabbit primary antibody, and Goat Anti-Mouse/Rabbit IgG (H&L)-HRP polymer. The section was then counterstained with haematoxylin and mounted with Neutral Gum.



IHC (Immunohistochemistry)

Immunohistochemical analysis staining in human liver carcinoma formalin fixed paraffin-embedded tissue section. The section was pre-treated using pressure cooker heat antigen retrieval with sodium citrate buffer (0.01M, pH=6) for 3 minutes. The section was detected using mouse primary antibody, and Goat Anti-Mouse/Rabbit IgG (H&L)-HRP polymer. The section was then counterstained with haematoxylin and mounted with Neutral Gum.

ADDITIONAL INFORMATION

Related Product Information for IgG secondary antibody	Goat Polyclonal Secondary Antibody to Mouse/Rabbit IgG (H&L) HRP polymer labeled
Product Categories/Family for IgG secondary antibody	Secondary Antibodies

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/igg-secondary-antibody-immunohistochemistry-western-blot-elisa/266236>