

LDHA recombinant antibody

Recombinant Anti-LDHA Rabbit mAb

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA266261	Rabbit	Human, Mouse, Rat	Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

LDHA; Monoclonal Recombinant Antibody; Recombinant Anti-LDHA Rabbit mAb; L-lactate dehydrogenase A chain; LDH-A; Cell proliferation-inducing gene 19 protein; LDH muscle subunit; LDH-M; Renal carcinoma antigen NY-REN-59; LDHA recombinant antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Monoclonal
Specificity	Recognizes endogenous levels of LDHA protein
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in PBS, pH7.4, containing 50% glycerol, 0.2% BSA and 0.01% sodium azide.
Applicable Applications for LDHA recombinant antibody	ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within human LDHA. The exact sequence is proprietary.
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C for one year. Avoid freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of LDHA staining in MCF7 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 degree C in a humidified chamber. Cells were washed with PBST and incubated with an AF488-conjugated secondary antibody (green) in PBS at room temperature in the dark. Phalloidin - AF594 was used to stain Actin filaments (red). DAPI was used to stain the cell nuclei (blue).



IHC (Immunohistochemistry)

Immunohistochemical analysis of LDHA staining in rat kidney formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. Tyramide-AF488 (green) was used as the chromogen. The section was then counterstained with DAPI (blue).



WB (Western Blot)

Western blot analysis of LDHA expression in HEK293T (A), HeLa (B), Jurkat (C), mouse kidney (D), rat liver (E), rat kidney (F) whole cell lysates. (Predicted band size: 36 kD; Observed band size: 37 kD)

ADDITIONAL INFORMATION

Related Product Information for LDHA recombinant antibody	Recombinant rabbit monoclonal antibody to LDHA
Product Categories/Family for LDHA recombinant antibody	Primary Antibody

NCBI AND UNIPROT INFORMATION

NCBI GI #	207028494
NCBI GeneID	3939
NCBI Accession #	NP_001128711.1
NCBI GenBank Nucleotide #	NM_001135239.1
UniProt Accession #	P00338; P06151; P04642
Molecular Weight	332
NCBI Official Full Name	L-lactate dehydrogenase A chain isoform 2
NCBI Official Synonym Full Names	lactate dehydrogenase A
NCBI Official Symbol	LDHA
NCBI Official Synonym Symbols	LDH1; LDHM; GSD11; PIG19
NCBI Protein Information	L-lactate dehydrogenase A chain; LDH-A; LDH-M; LDH muscle subunit; lactate dehydrogenase M; proliferation-inducing gene 19; renal carcinoma antigen NY-REN-59; cell proliferation-inducing gene 19 protein
UniProt Protein Name	L-lactate dehydrogenase A chain
UniProt Gene Name	LDHA
UniProt Synonym Gene Names	LDH-A; LDH-M
UniProt Entry Name	LDHA_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/monoclonal/ldha-recombinant-antibody-immunocytochemistry-immunofluorescence-immunohistochemistry-western-blot-elisa/266261>