

Rabbit TEK Polyclonal Antibody | anti-TEK antibody

TEK

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA268473	Rabbit	Human, Mouse, Rat	ELISA, Immunohistochemistry, Immunofluorescence, Western Blot


PRODUCT SYNONYMS

TEK; Polyclonal Antibody; CD202B; GLC3E; TIE-2; TIE2; VMCM; VMCM1; anti-TEK antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Specificity	This antibody detects endogenous levels of TEK.
Purity/Purification	Affinity purification
Form/Format	Liquid in PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
Sequence Length	1124
Applicable Applications for anti-TEK antibody	ELISA, IHC (Immunohistochemistry), IF (Immunofluorescence), WB (Western Blot)
Immunogen	Recombinant protein of human TEK
Conjugation	APC
Cellular Localization	Cell membrane ; Single-pass type I membrane protein. Cell junction . Cell junction, focal adhesion . Cytoplasm, cytoskeleton. Secreted . Recruited to cell-cell contacts in quiescent endothelial cells (PubMed:18425120, PubMed:18425119). Colocalizes with the actin cytoskeleton and at actin stress fibers during cell spreading. Recruited to the lower surface of migrating cells, especially the rear end of the cell. Proteolytic processing gives rise to a soluble extracellular domain that is secreted (PubMed:11806244). Tissue specificity:Detected in umbilical vein endothelial cells. Proteolytic processing gives rise to a soluble extracellular domain that is detected in blood plasma (at protein level). Predominantly expressed in endothelial cells and their progenitors, the angioblasts. Has been directly found in placenta and lung, with a lower level in umbilical vein endothelial cells, brain and kidney.
Function	Catalytic activity:ATP + a [protein]-L-tyrosine = ADP + a [protein]-L-tyrosine phosphate.,Disease:Defects in TEK are a cause of dominantly inherited venous malformations (VMCM) [MIM:600195]; an error of vascular morphogenesis characterized by dilated, serpiginous channels.,
Function	This protein is a protein tyrosine-kinase transmembrane receptor for angiopoietin 1. It may constitute the earliest mammalian endothelial cell lineage marker. Probably regulates endothelial cell proliferation, differentiation and guides the proper patterning of endothelial cells during blood vessel formation.,similarity:Belongs to the protein kinase superfamily. Tyr protein kinase family.,similarity:Belongs to the protein kinase superfamily. Tyr protein kinase family. Tie subfamily.,similarity:Contains 1 protein kinase domain.,similarity:Contains 2 Ig-like C2-type (immunoglobulin-like) domains.,similarity:Contains 3 EGF-like domains.,similarity:Contains 3 fibronectin type-III domains.,tissue specificity:Predominantly expressed in endothelial cells and their progenitors, the angioblasts. Has been directly found in placenta and lung, with a lower level in umbilical vein endothelial cells, brain and kidney
Preparation and Storage	Store at -20°C for 1 year. Avoid freeze / thaw cycles

IMAGES



IHC (Immunohistochemistry)
Immunohistochemical analysis of paraffin- embedded Human stomach. 1,Antibody was diluted at 1:400 (4° overnight). 2, High-pressure and temperature EDTA, pH 8.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200 (room temperature, 30min).



WB (Western Blot)
Western blot analysis of CACO2 lysate, antibody was diluted at 1000. Secondary antibody was diluted at 1:20000

NCBI AND UNIPROT INFORMATION

NCBI GI #	88758596
NCBI GeneID	7010
NCBI Accession #	NP_000450.2
NCBI GenBank Nucleotide #	NM_000459.4
UniProt Accession #	Q02763; Q5TCU2; Q8IV34; A8K6W0; B4DH20; B4DHD3; D3DRK5; E7EWI2
Molecular Weight	120kD (Observed)
NCBI Official Full Name	angiopoietin-1 receptor isoform 1
NCBI Official Synonym Full Names	TEK receptor tyrosine kinase
NCBI Official Symbol	TEK
NCBI Official Synonym Symbols	TIE2; VMCM; GLC3E; TIE-2; VMCM1; CD202B
NCBI Protein Information	angiopoietin-1 receptor
UniProt Protein Name	Angiopoietin-1 receptor
UniProt Gene Name	TEK
UniProt Synonym Gene Names	TIE2; VMCM; VMCM1; hTIE2

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/tek-antibody-immunohistochemistry-immunofluorescence-western-blot-elisa/268473>