

COVID 19 Spike S1 Coronavirus Active Protein | COVID-19 active protein

SARS-CoV-2 (2019-nCoV) Spike Protein (S1 Subunit, His Tag)

CATALOG #
AAA268893

HOST
HEK293 CellsSource: 2019nCoV


PRODUCT SYNONYMS

COVID 19 Spike S1 Coronavirus; SARS-CoV-2 (2019-nCoV) Spike Protein (S1 Subunit; His Tag); 2019 Novel Coronavirus; Coronavirus; CoV; COVID-19 virus; HCoV-2; Human Coronavirus 2019; SARS2; SARS-CoV-2; Severe acute respiratory syndrome coronavirus 2; Spike S1; COVID-19 active protein


PRODUCT SPECIFICATIONS

Host	HEK293 Cells Source: 2019nCoV
Purity/Purification	>90 % as determined by SDS-PAGE
Form/Format	Liquid or lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization.
Predicted N terminal	Val 16
Biological Activity	Measured by its binding ability in a functional ELISA. Immobilized human ACE2 protein (Ftag) at 10ug/mL (100uL/well) can bind 2019-nCoV Spike Protein (S1, His tag) (ZD2019-P7), the EC ₅₀ of 2019-nCoV Spike Protein (S1, His tag) (ZD2019-P7) is 200-400 ng/mL.
Protein Construction	A DNA sequence encoding the SARS-CoV-2 (2019-nCoV) spike protein S1Subunit (YP_009724390.1) (Val16-Arg685) was expressed with a polyhistidine tag at the C-terminus
Molecular Mass	The recombinant SARS-CoV-2 (2019-nCoV) Spike Protein (S1 Subunit, His Tag) consists of 681 amino acids and predicts a molecular mass of 76.5 kDa.
Dry Ice Note	The product may be shipped with dry ice, and an additional fee may be added to your shipping cost.
Preparation and Storage	Samples are stable for twelve months from date of receipt at -20 degree C to -80 degree C. Store it under sterile conditions at -20 degree C to -80 degree C upon receiving. Recommend to aliquot the protein into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles.

IMAGES



Application Data





Application Data

ADDITIONAL INFORMATION

Related Product Information for COVID-19 active protein	Description: The spike (S) glycoprotein of coronaviruses contains protrusions that will only bind to certain receptors on the host cell. Known receptors bind S1 are ACE2, angiotensin-converting enzyme2; DPP4, dipeptidyl peptidase-4; APN, aminopeptidase N; CEACAM, carcinoembryonic antigen-related cell adhesion molecule 1; Sia, sialic acid; O-ac Sia, O-acetylated sialic acid. The spike is essential for both host specificity and viral infectivity. The term 'peplomer' is typically used to refer to a grouping of heterologous proteins on the virus surface that function together. The spike (S) glycoprotein of coronaviruses is known to be essential in the binding of the virus to the host cell at the advent of the infection process.
References	References: 1. Shen S, et al. (2007) Expression, glycosylation, and modification of the spike (S) glycoprotein of SARS CoV. Methods Mol Biol. 379: 127-35. 2. Du L, et al. (2009) The spike protein of SARS-CoV--a target for vaccine and therapeutic development. Nat Rev Microbiol. 7 (3): 226-36. 3. Xiao X, et al. (2004) The SARS-CoV S glycoprotein. Cell Mol Life Sci. 61 (19-20): 2428-30

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/hek293-cells-source-2019ncov-covid-19-active-protein/covid-19-spike-s1-coronavirus/268893>