

Rabbit ICAM5 Polyclonal Antibody | anti-ICAM5 antibody

ICAM5

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA268998	Rabbit	Human, Mouse, Rat, Pig, Cow	Flow Cytometry, Functional Assay, Western Blot


PRODUCT SYNONYMS

ICAM5; Polyclonal Antibody; ICAM5_HUMAN; ICAM5 precursor; Intercellular adhesion molecule 5; Intercellular adhesion molecule 5 precursor; telencephalin; Telencephalin; TLCN; TLN; anti-ICAM5 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat, Pig, Cow
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Affinity purified by Protein A
Form/Format	0.01M TBS (pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.
Applicable Applications for anti-ICAM5 antibody	FCM/FACS (Flow Cytometry), WB (Western Blot)
Immunogen	KLH conjugated synthetic peptide derived from human ICAM5 501-600/924
Antibody type	Rabbit polyclonal Antibody
Preparation and Storage	Store at +4 degree C short term (1-2 weeks). Upon delivery aliquot. Store at -20 degree C. Avoid freeze / thaw cycle. Store In the Dark.

IMAGES



FCM/FACS (Flow Cytometry)

Blank control: HeLa.Primary Antibody (green line): Rabbit Anti-ICAM5 antibody.Dilution: 1ug/106 cells; Isotype Control Antibody (orange line): Rabbit IgG .Secondary Antibody : Goat anti-rabbit IgG-PE.Dilution: 0.2ug /test.Protocol: The cells were incubated in 5 %BSA to block non-specific protein-protein interactions for 30 min at at room temperature. Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.



WB (Western Blot)

Sample: Hippocampus (Mouse) Lysate at 40 ug.Primary: Anti- ICAM5 (AAA268998) at 1/1000 dilution.Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution.Predicted band size: 97 kD.Observed band size: 167 kD.

NCBI AND UNIPROT INFORMATION

UniProt Accession #	Q9UMF0
---------------------	--------

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/icam5-antibody-flow-cytometry-western-blot-elisa/268998>