

Rabbit iNOS Polyclonal Antibody | anti-iNOS antibody

Anti-iNOS Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA269599	Rabbit	Human, Mouse, Rat	Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

iNOS; Polyclonal Antibody; Anti-iNOS Antibody; NOS2A; Nitric oxide synthase; inducible; Hepatocyte NOS; HEP-NOS; Inducible NO synthase; Inducible NOS; NOS type II; Peptidyl-cysteine S-nitrosylase NOS2; anti-iNOS antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Applicable Applications for anti-iNOS antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human iNOS. The exact sequence is proprietary.
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C for one year. Avoid freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of iNOS staining in A549 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 degree C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).



IHC (Immunohistochemistry)

Immunohistochemical analysis of iNOS staining in human lung cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of iNOS expression in MCF7 (A), A549 (B), Raw264.7 (C), PC12 (D), rat liver (E) whole cell lysates.

ADDITIONAL INFORMATION

Related Product Information for anti-iNOS antibody	Rabbit polyclonal antibody to iNOS
--	------------------------------------

NCBI AND UNIPROT INFORMATION

NCBI GI #	24041029
NCBI GeneID	4843
NCBI Accession #	NP_000616.3
NCBI GenBank Nucleotide #	NM_000625.4
UniProt Accession #	P35228; O60757; O94994; Q16263; Q16692; Q4TTS5; Q9UD42; A1L3U5; B7ZLY2
Molecular Weight	131,117 Da
NCBI Official Full Name	nitric oxide synthase, inducible
NCBI Official Synonym Full Names	nitric oxide synthase 2, inducible
NCBI Official Symbol	NOS2
NCBI Official Synonym Symbols	NOS; INOS; NOS2A; HEP-NOS
NCBI Protein Information	nitric oxide synthase, inducible; NOS type II; NOS, type II; inducible NOS; hepatocyte NOS; inducible NO synthase; nitric oxide synthase, macrophage; peptidyl-cysteine S-nitrosylase NOS2; nitric oxide synthase 2A (inducible, hepatocytes)
UniProt Protein Name	Nitric oxide synthase, inducible
UniProt Gene Name	NOS2
UniProt Synonym Gene Names	NOS2A; HEP-NOS; Inducible NOS; INOS
UniProt Entry Name	NOS2_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.