

Rabbit TPL2 Polyclonal Antibody | anti-TPL2 antibody

Anti-TPL2 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA269664	Rabbit	Human, Mouse, Rat, Bovine, Monkey	Immunofluorescence, Immunohistochemistry, Western Blot

PRODUCT SYNONYMS

TPL2; Polyclonal Antibody; Anti-TPL2 Antibody; COT; ESTF; Mitogen-activated protein kinase kinase kinase 8; Cancer Osaka thyroid oncogene; Proto-oncogene c-Cot; Serine/threonine-protein kinase cot; Tumor progression locus 2; TPL-2; anti-TPL2 antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat, Bovine, Monkey
Clonality	Polyclonal
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Applicable Applications for anti-TPL2 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human TPL2. The exact sequence is proprietary.
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C for one year. Avoid freeze/thaw cycles.

IMAGES



IF (Immunofluorescence)

Immunofluorescent analysis of TPL2 staining in HEK293T cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 degree C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).



IHC (Immunohistochemistry)

Immunohistochemical analysis of TPL2 staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of TPL2 expression in SGC7901 (A), H9C2 (B), AML12 (C) whole cell lysates.

ADDITIONAL INFORMATION

Related Product Information for anti-TPL2 antibody	Rabbit polyclonal antibody to TPL2
--	------------------------------------

Product Categories/Family for anti-TPL2 antibody

Primary Antibody

NCBI AND UNIPROT INFORMATION

NCBI GI #	346644795
NCBI GeneID	1326
NCBI Accession #	NP_001231063.1
NCBI GenBank Nucleotide #	NM_001244134.1
UniProt Accession #	P41279; Q14275; Q5T855; Q9HC81; A8K2Q5; D3DRX1
Molecular Weight	52,925 Da
NCBI Official Full Name	mitogen-activated protein kinase kinase kinase 8
NCBI Official Synonym Full Names	mitogen-activated protein kinase kinase kinase 8
NCBI Official Symbol	MAP3K8
NCBI Official Synonym Symbols	COT; EST; ESTF; TPL2; MEKK8; Tpl-2; c-COT
NCBI Protein Information	mitogen-activated protein kinase kinase kinase 8; proto-oncogene c-Cot; tumor progression locus 2; Ewing sarcoma transformant; cot (cancer Osaka thyroid) oncogene; proto-oncogene serine/threonine protein kinase
UniProt Protein Name	Mitogen-activated protein kinase kinase kinase 8
UniProt Gene Name	MAP3K8
UniProt Synonym Gene Names	COT; ESTF; TPL-2
UniProt Entry Name	M3K8_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/tpl2-antibody-immunofluorescence-immunohistochemistry-western-blot-elisa/269664>