

Mouse OCT4 Monoclonal Antibody | anti-OCT4 antibody

OCT4 Monoclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA27015	Mouse	Human, Mouse, Rat	ELISA, Western Blot, Immunohistochemistry, Immunofluorescence, Flow Cytometry, Functional Assay


PRODUCT SYNONYMS

OCT4; Monoclonal Antibody; OCT4 Monoclonal Antibody; POU domain; class 5; transcription factor 1; Octamer-binding protein 3; Oct-3; Octamer-binding protein 4; Oct-4; Octamer-binding transcription factor 3; OTF-3; POU5F1; OCT3 OCT4 OTF3; anti-OCT4 antibody

PRODUCT SPECIFICATIONS


Host	Mouse
Reactivity	Human, Mouse, Rat
Clonality	Monoclonal
Isotype	IgG2b
Purity/Purification	>95% Protein G Purified
Form/Format	Liquid
Sequence Length	265
Applicable Applications for anti-OCT4 antibody	ELISA, WB (Western Blot), IHC (Immunohistochemistry), IF (Immunofluorescence), FCM/FACS (Flow Cytometry)
Application Notes	WB: 1:500-1:10000 IHC: 1:50-1:200 IF: 1:50-1:200
Immunogen	Recombinant Human POU domain, class 5, transcription factor 1 protein (1-360AA)
Conjugate	Non-Conjugated
Preservative	0.03% Proclin 300
Constituents	50% Glycerol, 0.01M PBS, pH 7.4
Preparation and Storage	Upon receipt, store at -20 degree C or -80 degree C. Avoid repeated freeze.

IMAGES



FCM (Flow Cytometry)

Overlay histogram showing Ntera-2 cells stained with CSB-MA018403A0m (red line) at 1:100. The cells were incubated in 1x PBS /10% normal goat serum to block non-specific protein-protein interactions followed by primary antibody for 1 h at 4 degree C. The secondary antibody used was FITC goat anti-mouse IgG(H+L) at 1/200 dilution for 1 h at 4 degree C. Isotype control antibody (green line) was used under the same conditions. Acquisition of >10,000 events was performed.



FCM (Flow Cytometry)

Overlay histogram showing A549 cells stained with CSB-MA018403A0m (red line) at 1:100. The cells were incubated in 1x PBS /10% normal goat serum to block non-specific protein-protein interactions followed by primary antibody for 1 h at 4 degree C. The secondary antibody used was FITC goat anti-mouse IgG(H+L) at 1/200 dilution for 1 h at 4 degree C. Isotype control antibody (green line) was used under the same conditions. Acquisition of >10,000 events was performed.




IF (Immunofluorescence)

Immunofluorescence staining of Ntera-2 cells with CSB-MA018403A0m at 1:100, counter-stained with DAPI. The cells were blocked in 10% normal Goat Serum and then incubated with the primary antibody overnight at 4 degree C. The secondary antibody was Alexa Fluor 488-conjugated AffiniPure Goat Anti-Mouse IgG(H+L).




IF (Immunofluorescence)

Immunofluorescence staining of A549 cells with CSB-MA018403A0m at 1:100, counter-stained with DAPI. The cells were blocked in 10% normal Goat Serum and then incubated with the primary antibody overnight at 4 degree C. The secondary antibody was Alexa Fluor 488-conjugated AffiniPure Goat Anti-Mouse IgG(H+L).



IHC (Immunohistochemistry)

IHC image of CSB-MA018403A0m diluted at 1:100 and staining in paraffin-embedded human endometrial cancer performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



IHC (Immunohistochemistry)

IHC image of CSB-MA018403A0m diluted at 1:100 and staining in paraffin-embedded human colon cancer performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.

ADDITIONAL INFORMATION

Product Categories/Family for anti-OCT4 antibody

Epigenetics and Nuclear Signaling

NCBI AND UNIPROT INFORMATION

NCBI GI #	553727233
NCBI GeneID	5460
NCBI Accession #	NP_001272916.1
NCBI GenBank Nucleotide #	NM_001285987.1
UniProt Accession #	Q01860; P31359; Q15167; Q15168; Q16422; Q5STF3; Q5STF4; A6NCS1; A6NLL8; D2IYK4
Molecular Weight	30,085 Da
NCBI Official Full Name	POU domain, class 5, transcription factor 1 isoform 3
NCBI Official Synonym Full Names	POU class 5 homeobox 1
NCBI Official Symbol	POU5F1
NCBI Official Synonym Symbols	OCT3; OCT4; OTF3; OTF4; OTF-3; Oct-3; Oct-4
NCBI Protein Information	POU domain, class 5, transcription factor 1
UniProt Protein Name	POU domain, class 5, transcription factor 1
UniProt Gene Name	POU5F1
UniProt Synonym Gene Names	OCT3; OCT4; OTF3; Oct-3; Oct-4; OTF-3

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/monoclonal/oct4-antibody-western-blot-immunohistochemistry-immunofluorescence-flow-cytometry-elisa/27015>