

Human IGF2R (Insulin Like Growth Factor 2 Receptor) ELISA Kit | IGF2R elisa kit

Human IGF2R (Insulin Like Growth Factor 2 Receptor) ELISA Kit

CATALOG #
AAA270677

REACTIVITY
Human


PRODUCT SYNONYMS

IGF2R (Insulin Like Growth Factor 2 Receptor); Human IGF2R (Insulin Like Growth Factor 2 Receptor) ELISA Kit; CD222; IGF2-R; CIMPR; M6P-R; MPRI; Cation-Independent Mannose-6 Phosphate Receptor; M6P/IGF2 receptor; 300 kDa mannose 6-phosphate receptor; IGF2R elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	This assay has high sensitivity and excellent specificity for detection of Human IGF2R. No significant cross-reactivity or interference between Human IGF2R and analogues was observed.
Assay Type	Sandwich
Detection Range	0.63-40 ng/mL
Sensitivity	0.221 ng/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay)? CV%<8%. Three samples of known concentration were tested twenty times on one plate to assess intra-assay precision.
Inter-assay Precision	Inter-assay Precision (Precision between assays)? CV%<10%. Three samples of known concentration were tested in forty separate assays to assess inter-assay precision.
Preparation and Storage	Store at 2-8 degree C. Stable for 6 months from date of receipt.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for IGF2R elisa kit

Principle of the Assay: The test principle applied in this kit is Sandwich enzyme immunoassay. The microtiter plate provided in this kit has been pre-coated with an antibody specific to Insulin Like Growth Factor 2 Receptor(IGF2R). Standards or samples are added to the appropriate microtiter plate wells then with a biotin-conjugated antibody specific to Insulin Like Growth Factor 2 Receptor(IGF2R). Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain Insulin Like Growth Factor 2 Receptor(IGF2R), biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm ± 10nm. The concentration of Insulin Like Growth Factor 2 Receptor(IGF2R) in the samples is then determined by comparing the OD of the samples to the standard curve.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/human-elisa-kits/igf2r-insulin-like-growth-factor-2-receptor/270677>