

Mouse PDE1C (Phosphodiesterase 1C, Calmodulin Dependent) ELISA Kit | PDE1C elisa kit

Mouse PDE1C (Phosphodiesterase 1C, Calmodulin Dependent) ELISA Kit

CATALOG #
AAA275066

REACTIVITY
Mouse


PRODUCT SYNONYMS

PDE1C (Phosphodiesterase 1C; Calmodulin Dependent); Mouse PDE1C (Phosphodiesterase 1C; Calmodulin Dependent) ELISA Kit; Hcam3; Cam-PDE 1C; Calcium/calmodulin-dependent 3'; 5'-cyclic nucleotide phosphodiesterase 1C; PDE1C elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Mouse
Specificity	This assay has high sensitivity and excellent specificity for detection of Mouse PDE1C. No significant cross-reactivity or interference between Mouse PDE1C and analogues was observed.
Assay Type	Sandwich
Detection Range	0.16-10 ng/mL
Sensitivity	0.053 ng/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay)? CV%<8%. Three samples of known concentration were tested twenty times on one plate to assess intra-assay precision.
Inter-assay Precision	Inter-assay Precision (Precision between assays)? CV%<10%. Three samples of known concentration were tested in forty separate assays to assess inter-assay precision.
Preparation and Storage	Store at 2-8 degree C. Stable for 6 months from date of receipt.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for PDE1C elisa kit	Principle of the Assay: The test principle applied in this kit is Sandwich enzyme immunoassay. The microtiter plate provided in this kit has been pre-coated with an antibody specific to Phosphodiesterase 1C, Calmodulin Dependent(PDE1C). Standards or samples are added to the appropriate microtiter plate wells then with a biotin-conjugated antibody specific to Phosphodiesterase 1C, Calmodulin Dependent(PDE1C). Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain Phosphodiesterase 1C, Calmodulin Dependent(PDE1C), biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm ± 10nm. The concentration of Phosphodiesterase 1C, Calmodulin Dependent(PDE1C) in the samples is then determined by comparing the OD of the samples to the standard curve.
Product Categories/Family for PDE1C elisa kit	Enzyme & Kinase

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/mouse-elisa-kits/pde1c-phosphodiesterase-1c-calmodulin-dependent/275066>