

General LXB4 (Lipoxin B4) ELISA Kit | LXB4 elisa kit

Universal LXB4 (Lipoxin B4) ELISA Kit

CATALOG #
AAA27526

REACTIVITY
General


PRODUCT SYNONYMS

LXB4 (Lipoxin B4); Universal LXB4 (Lipoxin B4) ELISA Kit; LXB4 elisa kit

PRODUCT SPECIFICATIONS

Reactivity	General
Specificity	Specifically recognize LXB4, no obvious cross reaction with other analogues
Assay Type	Competitive
Samples	Serum, plasma, cell culture supernatant and other biological samples
Detection Range	7.813-500pg/ml
Sensitivity	4.688pg/ml
Intra-assay Precision	Intra-assay Precision: samples with low, medium and high concentration are tested 20 times on same plate.
Inter-assay Precision	Inter-assay Precision: samples with low, medium and high concentration are tested 20 times on three different plates.
Preparation and Storage	Store entire kit at 2-8C for short-term. For longer-term, please store the microplate & standard at -20C, while the remaining reagents can be stored at 2-8C

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for LXB4 elisa kit	Background: Lipoxin B 4 (LXB 4) is a member of the lipoxin family, with its pro-resolving effects shown in allergic airway inflammation. In the upper airway, LXB4 significantly decreased nasal mucosal leukocytes and degranulation of mast cells (MCs) and eosinophils. In the lower airway, LXB4 significantly decreased airway inflammation, mucus metaplasia, and hyper-responsiveness. Principle of the Assay: This kit was based on Competitive-ELISA detection method. The microtiter plate provided in this kit has been pre-coated with LXB4. During the reaction, LXB4 in the sample or standard competes with a fixed amount of LXB4 on the solid phase supporter for sites on the Biotinylated Detection Antibody specific to LXB4. Excess conjugate and unbound sample or standard are washed from the plate, and HRP-Streptavidin (SABC) is added to each microplate well and incubated. Then TMB substrate solution is added to each well. The enzymesubstrate reaction is terminated by the addition of a acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm. The concentration of LXB4 in the samples is then determined by comparing the OD of the samples to the standard curve. The concentration of the target substance was inversely proportional to the OD450 value.
--	--

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/general-elisa-kits/lxb4-lipoxin-b4/27526>