

Mouse CAMK2gamma (Calcium/Calmodulin Dependent Protein Kinase II Gamma) ELISA Kit | CAMK2gamma elisa kit

Mouse CAMK2gamma (Calcium/Calmodulin Dependent Protein Kinase II Gamma) ELISA Kit

CATALOG # AAA277729	REACTIVITY Mouse
-------------------------------	----------------------------


PRODUCT SYNONYMS

CAMK2gamma (Calcium/Calmodulin Dependent Protein Kinase II Gamma); Mouse CAMK2gamma (Calcium/Calmodulin Dependent Protein Kinase II Gamma) ELISA Kit; CAMKII; CAMK2G; CAMK; CAMK-II; CAMKG; CaM Kinase; CaM kinase II subunit gamma; CaMK-II subunit gamma; CAMK2gamma elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Mouse
Specificity	This assay has high sensitivity and excellent specificity for detection of Mouse CAMK2 gamma. No significant cross-reactivity or interference between Mouse CAMK2gamma and analogues was observed.
Assay Type	Sandwich
Detection Range	0.16-10 ng/mL
Sensitivity	0.078 ng/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay)? CV% < 8%. Three samples of known concentration were tested twenty times on 1 plate to assess intra-assay precision.
Inter-assay Precision	Inter-assay Precision (precision between assays)? CV% < 10%. Three samples of known concentration were tested in forty separate assays to assess inter-assay precision.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for CAMK2gamma elisa kit

Principle of the Assay: The test principle applied in this kit is Sandwich enzyme immunoassay. The microtiter plate provided in this kit has been pre-coated with an antibody specific to Mouse CAMK2gamma. Standards or samples are added to the appropriate microtiter plate wells then with a biotin-conjugated antibody specific to Mouse CAMK2gamma. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain Mouse CAMK2gamma, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm ± 10nm. The concentration of Mouse CAMK2gamma in the samples is then determined by comparing the OD of the samples to the standard curve.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/mouse-elisa-kits/camk2gamma-calcium-calmodulin-dependent-protein-kinase-ii-gamma/277729>