

Tumor necrosis factor receptor superfamily member 9 (TNFRSF9) Active Protein | TNFRSF9 active protein

Recombinant Human Tumor necrosis factor receptor superfamily member 9 (TNFRSF9), partial (Active)


CATALOG # AAA278914	HOST Mammalian Cell
-------------------------------	-------------------------------

PRODUCT SYNONYMS


Tumor necrosis factor receptor superfamily member 9 (TNFRSF9); Recombinant Human Tumor necrosis factor receptor superfamily member 9 (TNFRSF9); partial (Active); (4-1BB ligand receptor)(CDw137)(T-cell antigen 4-1BB homolog)(T-cell antigen ILA)(CD antigen CD137); TNFRSF9 active protein

PRODUCT SPECIFICATIONS

Host	Mammalian Cell
Purity/Purification	Greater or equal to 85% purity as determined by SDS-PAGE.
Form/Format	Lyophilized or liquid (Format to be determined during the manufacturing process)
Sequence Positions	24-186aa, Partial
Sequence	LQDPCSNCPA GTFCDNNRNQ ICSPCPPNSF SSAGGQRTCD ICRQCKGVFR TRKECSSTSN AECDCTPGFH CLGAGCSMCE QDCKQGQELT KKGCKDCCFG TFNDQKRGIC RPWTNCSLDG KSVLVNGTKE RDVVCGPSA DLSPGASSVT PPAPAREPGH SPQ
Species	Homo sapiens (Human)
Endotoxin	Less than 1.0EU/ug as determined by LAL method.
Relevance	Receptor for TNFSF9/4-1BBL. Possibly active during T cell activation.
Reconstitution	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water. We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20 degree C/-80 degree C. Our default final concentration of glycerol is 50%. Customers could use it as reference.
Biological Activity	Measured by its binding ability in a functional ELISA. Immobilized TNFRSF9 at 2ug/mL can bind TNFSF9, the EC50 is 1.011-2.429ng/mL.
Preparation and Storage	Store at -20 degree C. For extended storage, store at -20 or -80 degree C. Repeated freezing and thawing is not recommended. Store working aliquots at 4 degree C for up to one week.



Bioactivity



SDS-PAGE
Tris-Glycine gel) Discontinuous SDS-PAGE (reduced) with 5% enrichment gel and 15% separation gel.

ADDITIONAL INFORMATION

Product Categories/Family for TNFRSF9 active protein	Cancer
--	--------

NCBI AND UNIPROT INFORMATION

NCBI GI #	5730095
NCBI GeneID	3604
NCBI Accession #	NP_001552.2
NCBI GenBank Nucleotide #	NM_001561.5
UniProt Accession #	Q07011; AAB34924
Molecular Weight	27,899 Da
NCBI Official Full Name	tumor necrosis factor receptor superfamily member 9
NCBI Official Synonym Full Names	tumor necrosis factor receptor superfamily, member 9
NCBI Official Symbol	TNFRSF9
NCBI Official Synonym Symbols	ILA; 4-1BB; CD137; CDw137
NCBI Protein Information	tumor necrosis factor receptor superfamily member 9; CD137 antigen; T cell antigen ILA; T-cell antigen ILA; 4-1BB ligand receptor; homolog of mouse 4-1BB; receptor protein 4-1BB; T-cell antigen 4-1BB homolog; induced by lymphocyte activation (ILA); interl
UniProt Protein Name	Tumor necrosis factor receptor superfamily member 9
UniProt Gene Name	TNFRSF9
UniProt Synonym Gene Names	CD137; ILA
UniProt Entry Name	TNR9_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.