

OCN recombinant antibody

OCN Recombinant Monoclonal Antibody

CATALOG # AAA278958	HOST HEK293F Cell	REACTIVITY Human	APPLICATIONS Flow Cytometry, Functional Assay, Immunohistochemistry, ELISA
-------------------------------	-----------------------------	----------------------------	--

PRODUCT SYNONYMS

OCN; Monoclonal Recombinant Antibody; OCN Recombinant Monoclonal Antibody; Occludin; OCN recombinant antibody

PRODUCT SPECIFICATIONS

Host	HEK293F Cell
Reactivity	Human
Clonality	Monoclonal
Isotype	Rabbit IgG
Clone Number	8H8
Purity/Purification	Affinity purified
Form/Format	Liquid; PBS, 150mM NaCl, 0.02% sodium azide, 50% glycerol, pH 7.4.
Applicable Applications for OCN recombinant antibody	FCM/FACS (Flow Cytometry), IHC (Immunohistochemistry), ELISA
Immunogen Species	Human
Immunogen	A synthesized peptide derived from Human OCN
Conjugation	Non-conjugated
Research Area	Signal transduction
Preparation and Storage	Upon receipt, store at -20 degree C or -80 degree C. Avoid repeated freeze.

IMAGES



FCM/FACS (Flow Cytometry)

Overlay Peak curve showing HepG2 cells stained with AAA278958 (red line) at 1:100. The cells were incubated in 10% normal goat serum to block non-specific protein-protein interactions followed by the antibody (1ug/1*106cells) for 45min at 4?. The secondary antibody used was FITC-conjugated Goat Anti-rabbit IgG(H+L) at 1:200 dilution for 35min at 4?.Control antibody (green line) was rabbit IgG (1ug/1*106cells) used under the same conditions. Acquisition of >10,000 events was performed.



IHC (Immunohistochemistry)

IHC image of AAA278958 diluted at 1:100 and staining in paraffin-embedded human colon cancer performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a Goat anti-rabbit polymer IgG labeled by HRP and visualized using 0.05% DAB.

ADDITIONAL INFORMATION

Related Product Information for OCN recombinant antibody	May play a role in the formation and regulation of the tight junction (TJ) paracellular permeability barrier. It is able to induce adhesion when expressed in cells lacking tight junctions. {ECO:0000269 PubMed:19114660}.; (Microbial infection) Acts as a co-receptor for hepatitis C virus (HCV) in hepatocytes. {ECO:0000269 PubMed:19182773, ECO:0000269 PubMed:20375010}.
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	327478416
NCBI GeneID	100506658
NCBI Accession #	NP_001192184.1
NCBI GenBank Nucleotide #	NM_001205255.1
UniProt Accession #	Q16625; Q5U1V4; Q8N6K1; B5BU70; D2DU64; D2DU65; D2IGC0; D2IGC1; E2CYV9
Molecular Weight	59,144 Da
NCBI Official Full Name	occludin isoform b
NCBI Official Synonym Full Names	occludin
NCBI Official Symbol	OCLN
NCBI Official Synonym Symbols	BLCPMG
NCBI Protein Information	occludin; tight junction protein occludin
UniProt Protein Name	Occludin
UniProt Gene Name	OCLN
UniProt Entry Name	OCLN_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/monoclonal/ocln-recombinant-antibody-clone-8h8-flow-cytometry-immunohistochemistry-elisa/278958>