

TBK1 recombinant antibody

TBK1 Recombinant Monoclonal Antibody


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA278996	HEK293F Cell	Human	Flow Cytometry, Functional Assay, Immunohistochemistry, ELISA

PRODUCT SYNONYMS

TBK1; Monoclonal Recombinant Antibody; TBK1 Recombinant Monoclonal Antibody; Serine/threonine-protein kinase TBK1 (EC 2.7.11.1)(NF-kappa-B-activating kinase)(T2K)(TANK-binding kinase 1); NAK; TBK1 recombinant antibody


PRODUCT SPECIFICATIONS

Host	HEK293F Cell
Reactivity	Human
Clonality	Monoclonal
Isotype	Rabbit IgG
Clone Number	14D10
Purity/Purification	Affinity purified
Form/Format	Liquid; PBS, 150mM NaCl, 0.02% sodium azide, 50% glycerol, pH 7.4.
Applicable Applications for TBK1 recombinant antibody	FCM/FACS (Flow Cytometry), IHC (Immunohistochemistry), ELISA
Immunogen Species	Human
Immunogen	A synthesized peptide derived from Human TBK1
Conjugation	Non-conjugated
Research Area	Epigenetics and Nuclear Signaling; Signal transduction
Preparation and Storage	Upon receipt, store at -20 degree C or -80 degree C. Avoid repeated freeze.




FCM/FACS (Flow Cytometry)

Overlay Peak curve showing A549 cells stained with AAA278996 (red line) at 1:100. The cells were fixed in 4% formaldehyde and permeated by 0.2% TritonX-100. Then 10% normal goat serum to block non-specific protein-protein interactions followed by the antibody (1ug/1*10⁶cells) for 45min at 4°. The secondary antibody used was FITC-conjugated Goat Anti-rabbit IgG(H+L) at 1:200 dilution for 35min at 4°. Control antibody (green line) was rabbit IgG (1ug/1*10⁶cells) used under the same conditions. Acquisition of >10,000 events was performed.



IHC (Immunohistochemistry)

IHC image of AAA278996 diluted at 1:100 and staining in paraffin-embedded human liver cancer performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a Goat anti-rabbit polymer IgG labeled by HRP and visualized using 0.05% DAB.



IHC (Immunohistochemistry)

IHC image of AAA278996 diluted at 1:100 and staining in paraffin-embedded human testis tissue performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a Goat anti-rabbit polymer IgG labeled by HRP and visualized using 0.05% DAB.

ADDITIONAL INFORMATION

Related Product Information for TBK1 recombinant antibody	Serine/threonine kinase that plays an essential role in regulating inflammatory responses to foreign agents (PubMed:12692549, PubMed:14703513, PubMed:18583960, PubMed:12702806, PubMed:15367631, PubMed:10581243, PubMed:11839743, PubMed:15485837, PubMed:21138416, PubMed:25636800, PubMed:23453971, PubMed:23453972, PubMed:23746807, PubMed:26611359, PubMed:32404352). Following activation of toll-like receptors by viral or bacterial components, associates with TRAF3 and TANK and phosphorylates interferon regulatory factors (IRFs) IRF3 and IRF7 as well as DDX3X (PubMed:12692549, PubMed:14703513, PubMed:18583960, PubMed:12702806, PubMed:15367631, PubMed:25636800). This activity allows subsequent homodimerization and nuclear translocation of the IRFs leading to transcriptional activation of pro-inflammatory and antiviral genes including IFNA and IFNB (PubMed:12702806, PubMed:15367631, PubMed:25636800, PubMed:32972995). In order to establish such an antiviral state, TBK1 form several different complexes whose composition depends on the type of cell and cellular stimuli (PubMed:23453971, PubMed:23453972, PubMed:23746807). Plays a key role in IRF3 activation: acts by first phosphorylating innate adapter proteins MAVS, STING1 and TICAM1 on their pLxIS motif, leading to recruitment of IRF3, thereby licensing IRF3 for phosphorylation by TBK1 (PubMed:25636800, PubMed:30842653). Phosphorylated IRF3 dissociates from the adapter proteins, dimerizes, and then enters the nucleus to induce expression of interferons (PubMed:25636800). Thus, several scaffolding molecules including FADD, TRADD, MAVS, AZI2, TANK or TBKBP1/SINTBAD can be recruited to the TBK1-containing-complexes (PubMed:21931631). Under particular conditions, functions as a NF-kappa-B effector by phosphorylating NF-kappa-B inhibitor alpha/NFKBIA, IKBKB or RELA to translocate NF-Kappa-B to the nucleus (PubMed:10783893, PubMed:15489227). Restricts bacterial proliferation by phosphorylating the autophagy receptor OPTN/Optineurin on 'Ser-177', thus enhancing LC3 binding affinity and antibacterial autophagy (PubMed:21617041). Phosphorylates SMCR8 component of the C9orf72-SMCR8 complex, promoting autophagosome maturation (PubMed:27103069). Phosphorylates and activates AKT1 (PubMed:21464307). Seems to play a role in energy balance regulation by sustaining a state of chronic, low-grade inflammation in obesity, wich leads to a negative impact on insulin sensitivity (By similarity). Attenuates retroviral budding by phosphorylating the endosomal sorting complex required for transport-I (ESCRT-I) subunit VPS37C (PubMed:21270402). Phosphorylates Borna disease virus (BDV) P protein (PubMed:16155125). Plays an essential role in the TLR3- and IFN-dependent control of herpes virus HSV-1 and HSV-2 infections in the central nervous system (PubMed:22851595). {ECO:0000250 UniProtKB:Q9WUN2, ECO:0000269 PubMed:10581243, ECO:0000269 PubMed:10783893, ECO:0000269 PubMed:11839743, ECO:0000269 PubMed:12692549, ECO:0000269 PubMed:12702806, ECO:0000269 PubMed:14703513, ECO:0000269 PubMed:15367631, ECO:0000269 PubMed:15485837, ECO:0000269 PubMed:15489227, ECO:0000269 PubMed:16155125, ECO:0000269 PubMed:18583960, ECO:0000269 PubMed:21138416, ECO:0000269 PubMed:21270402, ECO:0000269 PubMed:21464307, ECO:0000269 PubMed:21617041, ECO:0000269 PubMed:21931631, ECO:0000269 PubMed:22851595, ECO:0000269 PubMed:23453971, ECO:0000269 PubMed:23453972, ECO:0000269 PubMed:23746807, ECO:0000269 PubMed:25636800, ECO:0000269 PubMed:26611359, ECO:0000269 PubMed:27103069, ECO:0000269 PubMed:30842653, ECO:0000269 PubMed:32972995}.
---	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	7019547
NCBI GeneID	29110
NCBI Accession #	NP_037386.1
NCBI GenBank Nucleotide #	NM_013254.3
UniProt Accession #	Q9UHD2; Q8IYV3; Q9NUJ5; A8K4S4
Molecular Weight	84kDa
NCBI Official Full Name	serine/threonine-protein kinase TBK1
NCBI Official Synonym Full Names	TANK-binding kinase 1
NCBI Official Symbol	TBK1
NCBI Official Synonym Symbols	NAK; T2K
NCBI Protein Information	serine/threonine-protein kinase TBK1; NF-kB-activating kinase; NF-kappa-B-activating kinase
UniProt Protein Name	Serine/threonine-protein kinase TBK1
UniProt Gene Name	TBK1

UniProt Synonym Gene Names	NAK
UniProt Entry Name	TBK1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/monoclonal/tbk1-recombinant-antibody-clone-14d10-flow-cytometry-immunohistochemistry-elisa/278996>