

CCR8 recombinant antibody

CCR8 Recombinant Monoclonal Antibody


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA279026	HEK293F Cell	Human	Flow Cytometry, Functional Assay

PRODUCT SYNONYMS


CCR8; Monoclonal Recombinant Antibody; CCR8 Recombinant Monoclonal Antibody; C C chemokine receptor type 8; C C CKR 8; C C motif chemokine receptor 8; C-C chemokine receptor type 8; C-C CKR-8; CC chemokine receptor 8; CC chemokine receptor CHEMR1; CC chemokine receptor type 8; CC CKR 8; CC-CKR-8; CCR 8; CCR-8; Ccr8; CCR8 protein; CCR8-L; CCR8_HUMAN; CDw198; CDw198 antigen; Chemokine (C C motif) receptor 8; Chemokine (C C) receptor 8; Chemokine (C C) receptor like 2; Chemokine (CC motif) receptor 8; Chemokine (CC) receptor 8; Chemokine (CC) receptor like 1; Chemokine (CC) receptor like 2; Chemokine C C motif receptor 8; Chemokine C C receptor 8; Chemokine CC motif receptor 8; Chemokine CC receptor 8; Chemokine receptor 8; Chemokine receptor like 1; Chemokine receptor-like 1; CKR L1; CKR-L1; CKRL 1; CKRL1; CMKBR 8; CMKBR L2; CMKBR8; CMKBRL 2; CMKBRL2; CY 6; CY6; GPR CY6; GPR-CY6; GPRCY6; MGC123958; MGC123959; MGC129966; MGC129973; TER 1; TER1; CCR8 recombinant antibody

PRODUCT SPECIFICATIONS

Host	HEK293F Cell
Reactivity	Human
Clonality	Monoclonal
Isotype	hIgG1, lambda 2
Clone Number	10A9
Purity/Purification	Affinity purified
Form/Format	Liquid; 0.03% Proclin 300, 50% glycerol, 0.01M PBS, pH 7.4.
Applicable Applications for CCR8 recombinant antibody	FCM/FACS (Flow Cytometry)
Immunogen Species	Human
Immunogen	Recombinant Human CCR8 protein
Conjugation	Non-conjugated
Research Area	Immunology
Preparation and Storage	Upon receipt, store at -20 degree C or -80 degree C. Avoid repeated freeze.



FCM/FACS (Flow Cytometry)
Untransfected CHO-K1 cells (green line) and transfected Human CCR8 CHO-K1 stable cells (red line) were stained with anti-CCR8 recombinant antibody (2ug/1*10⁶cells), washed and then followed by FITC-conjugated anti-Human IgG Fc antibody and analyzed with flow cytometry.



FCM/FACS (Flow Cytometry)
Untransfected HEK293T cells (green line) and transfected Human CCR8 HEK293T stable cells (red line) were stained with anti-CCR8 recombinant antibody (2ug/1*10⁶cells), washed and then followed by FITC-conjugated anti-Human IgG Fc antibody and analyzed with flow cytometry.

NCBI AND UNIPROT INFORMATION

NCBI GI #	1707884
NCBI GeneID	1237
UniProt Accession #	P51685; Q3KNQ8; Q3KNR3; Q9BYX5; B2RC64
NCBI Official Full Name	C-C chemokine receptor type 8
NCBI Official Synonym Full Names	chemokine (C-C motif) receptor 8
NCBI Official Symbol	CCR8
NCBI Official Synonym Symbols	CY6; TER1; CCR-8; CKRL1; CDw198; CMKBR8; GPRCY6; CMKBRL2; CC-CKR-8
NCBI Protein Information	C-C chemokine receptor type 8; CC chemokine receptor 8; chemokine receptor-like 1; chemokine (C-C) receptor 8; CC chemokine receptor CHEMR1; CC-chemokine receptor chemr1; chemokine (C-C) receptor-like 2
UniProt Protein Name	C-C chemokine receptor type 8
UniProt Gene Name	CCR8
UniProt Synonym Gene Names	CKRL1; CMKBR8; CMKBRL2; C-C CKR-8; CC-CKR-8; CCR-8; CKR-L1; GPRCY6
UniProt Entry Name	CCR8_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.