

Mouse anti-Human HIST1H3A Monoclonal Antibody | anti-HIST1H3A antibody

Mouse anti-Homo sapiens (Human) HIST1H3A Monoclonal Antibody antibody


CATALOG # AAA279038	HOST Mouse	REACTIVITY Human	APPLICATIONS Flow Cytometry, Functional Assay, ELISA
-------------------------------	----------------------	----------------------------	--

PRODUCT SYNONYMS


HIST1H3A; Monoclonal Antibody; Mouse anti-Homo sapiens (Human) HIST1H3A Monoclonal Antibody antibody; Phospho-Histone H3(Ser28) Monoclonal Antibody; Histone H3.1 (Histone H3/a)(Histone H3/b)(Histone H3/c)(Histone H3/d)(Histone H3/f)(Histone H3/h)(Histone H3/i)(Histone H3/j)(Histone H3/k)(Histone H3/l); HIST1H3B; HIST1H3C; HIST1H3D; HIST1H3E; HIST1H3F; HIST1H3G; HIST1H3H; HIST1H3I; HIST1H3J; H3FA; H3FL; H3FC; H3FB; H3FD; H3FI; H3FH; H3FK; H3FF; H3FJ; anti-HIST1H3A antibody

PRODUCT SPECIFICATIONS

Host	Mouse
Reactivity	Human
Clonality	Monoclonal
Isotype	IgG1
Clone Number	3F4C1
Purity/Purification	>95%, Protein G purified
Form/Format	Liquid; 0.03% Proclin 300, 50% glycerol, 0.01M PBS, pH 7.4.
Applicable Applications for anti-HIST1H3A antibody	FCM/FACS (Flow Cytometry), ELISA
Immunogen Species	Human
Immunogen	KS(Phospho)APATGGVKKPHRC
Conjugation	Non-conjugated
Research Area	Epigenetics and Nuclear Signaling
Preparation and Storage	Upon receipt, store at -20 degree C or -80 degree C. Avoid repeated freeze.



FCM/FACS (Flow Cytometry)
Overlay Peak curve showing HepG2 cells stained with AAA279038 (red line) at 1:100. The cells were incubated in 10% normal goat serum to block non-specific protein-protein interactions followed by the antibody (1ug/1*10⁶cells) for 1h at 4°C. The secondary antibody used was FITC-conjugated Goat Anti-Mouse IgG(H+L) at 1/100 dilution for 30min at 4°C. Isotype control antibody (green line) was mouse IgG1 (1ug/1*10⁶cells) used under the same conditions. Acquisition of >10,000 events was performed.



FCM/FACS (Flow Cytometry)
Overlay Peak curve showing Hela cells stained with AAA279038 (red line) at 1:100. The cells were incubated in 10% normal goat serum to block non-specific protein-protein interactions followed by the antibody (1ug/1*10⁶cells) for 1h at 4°C. The secondary antibody used was FITC-conjugated Goat Anti-Mouse IgG(H+L) at 1/100 dilution for 30min at 4°C. Isotype control antibody (green line) was mouse IgG1 (1ug/1*10⁶cells) used under the same conditions. Acquisition of >10,000 events was performed.

NCBI AND UNIPROT INFORMATION

NCBI GI #	4504281
NCBI GeneID	8350
NCBI Accession #	NP_003520.1
NCBI GenBank Nucleotide #	NM_003529.2
UniProt Accession #	P68431; P02295; P02296; P16106; Q6ISV8; Q6NWP8; Q6NWP9; Q6NXU4; Q71DJ3; Q93081; A0PJT7; A5PLR1
Molecular Weight	15,404 Da
NCBI Official Full Name	histone H3.1
NCBI Official Synonym Full Names	histone cluster 1, H3a
NCBI Official Symbol	HIST1H3A
NCBI Official Synonym Symbols	H3/A; H3FA
NCBI Protein Information	histone H3.1; histone H3/a; histone 1, H3a; H3 histone family, member A
UniProt Protein Name	Histone H3.1
UniProt Gene Name	HIST1H3A
UniProt Synonym Gene Names	H3FA
UniProt Entry Name	H31_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.