

RELB recombinant antibody

RELB Recombinant Monoclonal Antibody



CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA279089	HEK293F Cell	Human	Flow Cytometry, Functional Assay, Immunofluorescence, ELISA

PRODUCT SYNONYMS

RELB; Monoclonal Recombinant Antibody; RELB Recombinant Monoclonal Antibody; Transcription factor RelB (I-Rel); RELB recombinant antibody

PRODUCT SPECIFICATIONS

Host	HEK293F Cell
Reactivity	Human
Clonality	Monoclonal
Isotype	Rabbit IgG
Clone Number	9E4
Purity/Purification	Affinity purified
Form/Format	Liquid; PBS, 150mM NaCl, 0.02% sodium azide, 50% glycerol, pH 7.4.
Applicable Applications for RELB recombinant antibody	FCM/FACS (Flow Cytometry), IF (Immunofluorescence), ELISA
Immunogen Species	Human
Immunogen	A synthesized peptide derived from Human RELB
Conjugation	Non-conjugated
Research Area	Epigenetics and Nuclear Signaling; Cancer; Cell biology; Signal transduction
Preparation and Storage	Upon receipt, store at -20 degree C or -80 degree C. Avoid repeated freeze.



IF (Immunofluorescence)
Immunofluorescence staining of HeLa with AAA279089 at 1:25, counter-stained with DAPI. The cells were fixed in 4% formaldehyde and permeated by 0.2% TritonX-100. Then 10% normal goat serum to block non-specific protein-protein interactions followed by the antibody (1ug/1*10⁶cells) for 45min at 4?. The secondary antibody used was FITC-conjugated Goat Anti-rabbit IgG(H+L) at 1:200 dilution for 35min at 4?.Control antibody (green line) was rabbit IgG (1ug/1*10⁶cells) used under the same conditions. Acquisition of >10,000 events was performed.

FCM/FACS (Flow Cytometry)
Overlay Peak curve showing HeLa cells stained with AAA279089 (red line) at 1:50. The cells were fixed in 4% formaldehyde and permeated by 0.2% TritonX-100. Then 10% normal goat serum to block non-specific protein-protein interactions followed by the antibody (1ug/1*10⁶cells) for 45min at 4?. The secondary antibody used was FITC-conjugated Goat Anti-rabbit IgG(H+L) at 1:200 dilution for 35min at 4?.Control antibody (green line) was rabbit IgG (1ug/1*10⁶cells) used under the same conditions. Acquisition of >10,000 events was performed.

NCBI AND UNIPROT INFORMATION

NCBI GI #	35493878
NCBI GeneID	5971
NCBI Accession #	NP_006500.2
NCBI GenBank Nucleotide #	NM_006509.3
UniProt Accession #	Q01201; Q6GTX7; Q9UEI7
Molecular Weight	62,134 Da
NCBI Official Full Name	transcription factor RelB
NCBI Official Synonym Full Names	v-rel reticuloendotheliosis viral oncogene homolog B
NCBI Official Symbol	RELB
NCBI Official Synonym Symbols	IREL; I-REL; REL-B
NCBI Protein Information	transcription factor RelB; v-rel reticuloendotheliosis viral oncogene homolog B, nuclear factor of kappa light polypeptide gene enhancer in B-cells 3; v-rel avian reticuloendotheliosis viral oncogene homolog B (nuclear factor of kappa light polypeptide ge
UniProt Protein Name	Transcription factor RelB
UniProt Gene Name	RELB
UniProt Entry Name	RELB_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.