

SMARCA4 recombinant antibody

SMARCA4 Recombinant Monoclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA279097	HEK293F Cell	Human	Flow Cytometry, Functional Assay, Immunofluorescence, Immunohistochemistry, ELISA

PRODUCT SYNONYMS

SMARCA4; Monoclonal Recombinant Antibody; SMARCA4 Recombinant Monoclonal Antibody; Transcription activator BRG1 (EC 3.6.4.-)(ATP-dependent helicase SMARCA4)(BRG1-associated factor 190A)(BAF190A)(Mitotic growth and transcription activator)(Protein BRG-1)(Protein brahma homolog 1)(SNF2-beta)(SWI/SNF-related matrix-associated actin-dependent regulator of chromatin subfamily A member 4); BAF190A BRG1 SNF2B SNF2L4; SMARCA4 recombinant antibody

PRODUCT SPECIFICATIONS

Host	HEK293F Cell
Reactivity	Human
Clonality	Monoclonal
Isotype	Rabbit IgG
Clone Number	22F3
Purity/Purification	Affinity purified
Form/Format	Liquid; PBS, 150mM NaCl, 0.02% sodium azide, 50% glycerol, pH 7.4.
Applicable Applications for SMARCA4 recombinant antibody	FCM/FACS (Flow Cytometry), IF (Immunofluorescence), IHC (Immunohistochemistry), ELISA
Immunogen Species	Human
Immunogen	A synthesized peptide derived from Human SMARCA4
Conjugation	Non-conjugated
Research Area	Epigenetics and Nuclear Signaling; Neuroscience; Cancer; Stem cells
Preparation and Storage	Upon receipt, store at -20 degree C or -80 degree C. Avoid repeated freeze.

IMAGES



FCM/FACS (Flow Cytometry)

Overlay Peak curve showing Hela cells stained with AAA279097 (red line) at 1:50. The cells were fixed in 4% formaldehyde and permeated by 0.2% TritonX-100. Then 10% normal goat serum to block non-specific protein-protein interactions followed by the antibody (1ug/1*10⁶cells) for 45min at 4?. The secondary antibody used was FITC-conjugated Goat Anti-rabbit IgG(H+L) at 1:200 dilution for 35min at 4?. Control antibody (green line) was rabbit IgG (1ug/1*10⁶cells) used under the same conditions. Acquisition of >10,000 events was performed.



IF (Immunofluorescence)

Immunofluorescence staining of Hela with AAA279097 at 1:20, counter-stained with DAPI. The cells were fixed in 4% formaldehyde and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 511-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



IHC (Immunohistochemistry)

IHC image of AAA279097 diluted at 1:50 and staining in paraffin-embedded human breast cancer performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a Goat anti-rabbit polymer IgG labeled by HRP and visualized using 0.44% DAB.



IHC (Immunohistochemistry)

IHC image of AAA279097 diluted at 1:50 and staining in paraffin-embedded human stomach tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a Goat anti-rabbit polymer IgG labeled by HRP and visualized using 0.44% DAB.

NCBI AND UNIPROT INFORMATION

NCBI GI #	192807312
NCBI GeneID	6597
NCBI Accession #	NP_001122316.1
NCBI GenBank Nucleotide #	NM_001128844.1
UniProt Accession #	P51532; O95052; Q9HBD3; B1A8Z4; B1A8Z5; B1A8Z6; B1A8Z7; E9PBR8
Molecular Weight	1647
NCBI Official Full Name	transcription activator BRG1 isoform B
NCBI Official Synonym Full Names	SWI/SNF related, matrix associated, actin dependent regulator of chromatin, subfamily a, member 4
NCBI Official Symbol	SMARCA4
NCBI Official Synonym Symbols	BRG1; SNF2; SWI2; MRD16; RTPS2; BAF190; SNF2L4; SNF2LB; hSNF2b; BAF190A
NCBI Protein Information	transcription activator BRG1; SNF2-beta; SNF2-like 4; protein BRG-1; nuclear protein GRB1; brahma protein-like 1; BRM/SWI2-related gene 1; homeotic gene regulator; protein brahma homolog 1; BRG1-associated factor 190A; sucrose nonfermenting-like 4; ATP-de
UniProt Protein Name	Transcription activator BRG1
UniProt Gene Name	SMARCA4
UniProt Synonym Gene Names	BAF190A; BRG1; SNF2B; SNF2L4; BAF190A
UniProt Entry Name	SMCA4_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/monoclonal/smarca4-recombinant-antibody-clone-22f3-flow-cytometry-immunofluorescence-immunohistochemistry-elisa/279097>