

PRKAR2A recombinant antibody

PRKAR2A Recombinant Monoclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA279100	HEK293F Cell	Human	Flow Cytometry, Functional Assay, Immunofluorescence, Immunohistochemistry, ELISA


PRODUCT SYNONYMS

PRKAR2A; Monoclonal Recombinant Antibody; PRKAR2A Recombinant Monoclonal Antibody; cAMP-dependent protein kinase type II-alpha regulatory subunit; PKR2 PRKAR2; PRKAR2A recombinant antibody

PRODUCT SPECIFICATIONS


Host	HEK293F Cell
Reactivity	Human
Clonality	Monoclonal
Isotype	Rabbit IgG
Clone Number	15G8
Purity/Purification	Affinity purified
Form/Format	Liquid; PBS, 150mM NaCl, 0.02% sodium azide, 50% glycerol, pH 7.4.
Applicable Applications for PRKAR2A recombinant antibody	FCM/FACS (Flow Cytometry), IF (Immunofluorescence), IHC (Immunohistochemistry), ELISA
Immunogen Species	Human
Immunogen	A synthesized peptide derived from Human PRKAR2A
Conjugation	Non-conjugated
Research Area	Cancer; Metabolism; Signal transduction
Preparation and Storage	Upon receipt, store at -20 degree C or -80 degree C. Avoid repeated freeze.

IMAGES



FCM/FACS (Flow Cytometry)

Overlay Peak curve showing MCF-7 cells stained with AAA279100 (red line) at 1:50. The cells were fixed in 4% formaldehyde and permeated by 0.2% TritonX-100. Then 10% normal goat serum to block non-specific protein-protein interactions followed by the antibody (1ug/1*10⁶cells) for 45min at 4?. The secondary antibody used was FITC-conjugated Goat Anti-rabbit IgG(H+L) at 1:200 dilution for 35min at 4?.Control antibody (green line) was rabbit IgG (1ug/1*10⁶cells) used under the same conditions. Acquisition of >10,000 events was performed.




IF (Immunofluorescence)

Immunofluorescence staining of HeLa with AAA279100 at 1:5, counter-stained with DAPI. The cells were fixed in 4% formaldehyde and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 512-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



IHC (Immunohistochemistry)

IHC image of AAA279100 diluted at 1:50 and staining in paraffin-embedded human rectal cancer performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a Goat anti-rabbit polymer IgG labeled by HRP and visualized using 0.47% DAB.



IHC (Immunohistochemistry)

IHC image of AAA279100 diluted at 1:50 and staining in paraffin-embedded human kidney tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a Goat anti-rabbit polymer IgG labeled by HRP and visualized using 0.47% DAB.

NCBI AND UNIPROT INFORMATION

NCBI GI #	4758958
NCBI GeneID	5576
NCBI Accession #	NP_004148.1
NCBI GenBank Nucleotide #	NM_004157.2
UniProt Accession #	P13861; Q16823
Molecular Weight	45,518 Da
NCBI Official Full Name	cAMP-dependent protein kinase type II-alpha regulatory subunit
NCBI Official Synonym Full Names	protein kinase, cAMP-dependent, regulatory, type II, alpha
NCBI Official Symbol	PRKAR2A
NCBI Official Synonym Symbols	PKR2; PRKAR2
NCBI Protein Information	cAMP-dependent protein kinase type II-alpha regulatory subunit; protein kinase A, RII-alpha subunit; cAMP-dependent protein kinase regulatory subunit RII alpha
UniProt Protein Name	cAMP-dependent protein kinase type II-alpha regulatory subunit
UniProt Gene Name	PRKAR2A
UniProt Synonym Gene Names	PKR2; PRKAR2
UniProt Entry Name	KAP2_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/monoclonal/prkar2a-recombinant-antibody-clone-15g8-flow-cytometry-immunofluorescence-immunohistochemistry-elisa/279100>