

BAX recombinant antibody

BAX Recombinant Monoclonal Antibody



CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA279106	HEK293F Cell	Human	Flow Cytometry, Functional Assay, Immunohistochemistry, ELISA

PRODUCT SYNONYMS

BAX; Monoclonal Recombinant Antibody; BAX Recombinant Monoclonal Antibody; Apoptosis regulator BAX (Bcl-2-like protein 4)(Bcl2-L-4); BCL2L4; BAX recombinant antibody


PRODUCT SPECIFICATIONS

Host	HEK293F Cell
Reactivity	Human
Clonality	Monoclonal
Isotype	Rabbit IgG
Clone Number	15D9
Purity/Purification	Affinity purified
Form/Format	Liquid; PBS, 150mM NaCl, 0.02% sodium azide, 50% glycerol, pH 7.4.
Applicable Applications for BAX recombinant antibody	FCM/FACS (Flow Cytometry), IHC (Immunohistochemistry), ELISA
Immunogen Species	Human
Immunogen	A synthesized peptide derived from Human BAX
Conjugation	Non-conjugated
Research Area	Cancer; Cell biology; Metabolism
Preparation and Storage	Upon receipt, store at -20 degree C or -80 degree C. Avoid repeated freeze.



IHC (Immunohistochemistry)
IHC image of AAA279106 diluted at 1:50 and staining in paraffin-embedded human rectal cancer performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a Goat anti-rabbit polymer IgG labeled by HRP and visualized using 0.53% DAB.

FCM/FACS (Flow Cytometry)
Overlay Peak curve showing Hela cells stained with AAA279106 (red line) at 1:50. The cells were fixed in 4% formaldehyde and permeated by 0.2% TritonX-100. Then 10% normal goat serum to block non-specific protein-protein interactions followed by the antibody (1ug/1*10⁶cells) for 45min at 4°. The secondary antibody used was FITC-conjugated Goat Anti-rabbit IgG(H+L) at 1:200 dilution for 35min at 4°. Control antibody (green line) was rabbit IgG (1ug/1*10⁶cells) used under the same conditions. Acquisition of >10,000 events was performed.



IHC (Immunohistochemistry)
IHC image of AAA279106 diluted at 1:50 and staining in paraffin-embedded mouse lung tissue performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a Goat anti-rabbit polymer IgG labeled by HRP and visualized using 0.53% DAB.

NCBI AND UNIPROT INFORMATION

NCBI GI #	4757838
NCBI GeneID	581
NCBI Accession #	NP_004315.1
NCBI GenBank Nucleotide #	NM_004324.3
UniProt Accession #	Q07812; P55269; Q07814; Q07815; Q8WZ49; Q9NR76; Q9NYG7; Q9UCZ6; Q9UCZ7; Q9UQD6; A8K4W1
Molecular Weight	21,184 Da
NCBI Official Full Name	apoptosis regulator BAX isoform beta
NCBI Official Synonym Full Names	BCL2-associated X protein
NCBI Official Symbol	BAX
NCBI Official Synonym Symbols	BCL2L4
NCBI Protein Information	apoptosis regulator BAX; bcl2-L-4; bcl-2-like protein 4; BCL2-associated X protein omega; BCL2-associated X protein transcript variant delta2

UniProt Protein Name	Apoptosis regulator BAX
UniProt Gene Name	BAX
UniProt Synonym Gene Names	BCL2L4; Bcl2-L-4
UniProt Entry Name	BAX_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/monoclonal/bax-recombinant-antibody-clone-15d9-flow-cytometry-immunohistochemistry-elisa/279106>