

MDM2 recombinant antibody

Phospho-MDM2 (S166) Recombinant Monoclonal Antibody


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA279110	HEK293F Cell	Human	Flow Cytometry, Functional Assay, Immunofluorescence, Immunohistochemistry, ELISA

PRODUCT SYNONYMS

MDM2; Monoclonal Recombinant Antibody; Phospho-MDM2 (S166) Recombinant Monoclonal Antibody; E3 ubiquitin-protein ligase Mdm2 (EC 2.3.2.27)(Double minute 2 protein)(Hdm2)(Oncoprotein Mdm2)(RING-type E3 ubiquitin transferase Mdm2)(p53-binding protein Mdm2); MDM2 recombinant antibody


PRODUCT SPECIFICATIONS

Host	HEK293F Cell
Reactivity	Human
Clonality	Monoclonal
Isotype	Rabbit IgG
Clone Number	14F8
Purity/Purification	Affinity purified
Form/Format	Liquid; PBS, 150mM NaCl, 0.02% sodium azide, 50% glycerol, pH 7.4.
Applicable Applications for MDM2 recombinant antibody	FCM/FACS (Flow Cytometry), IF (Immunofluorescence), IHC (Immunohistochemistry), ELISA
Immunogen Species	Human
Immunogen	A synthesized peptide derived from Human MDM2
Conjugation	Non-conjugated
Research Area	Epigenetics and Nuclear Signaling; Cancer; Cell biology
Preparation and Storage	Upon receipt, store at -20 degree C or -80 degree C. Avoid repeated freeze.




FCM/FACS (Flow Cytometry)

Overlay Peak curve showing Hela cells stained with AAA279110 (red line) at 1:50. The cells were fixed in 4% formaldehyde and permeated by 0.2% TritonX-100. Then 10% normal goat serum to block non-specific protein-protein interactions followed by the antibody (1ug/1*106cells) for 45min at 4?. The secondary antibody used was FITC-conjugated Goat Anti-rabbit IgG(H+L) at 1:200 dilution for 35min at 4?.Control antibody (green line) was rabbit IgG (1ug/1*106cells) used under the same conditions. Acquisition of >10,000 events was performed.




IHC (Immunohistochemistry)

IHC image of AAA279110 diluted at 1:50 and staining in paraffin-embedded human lung cancer performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a Goat anti-rabbit polymer IgG labeled by HRP and visualized using 0.57% DAB.



IF (Immunofluorescence)

Immunofluorescence staining of HepG2 with AAA279110 at 1:10, counter-stained with DAPI. The cells were fixed in 4% formaldehyde and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 517-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



IHC (Immunohistochemistry)

IHC image of AAA279110 diluted at 1:50 and staining in paraffin-embedded human brain tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a Goat anti-rabbit polymer IgG labeled by HRP and visualized using 0.57% DAB.

NCBI AND UNIPROT INFORMATION

NCBI GI #	89993689
NCBI GeneID	4193
NCBI Accession #	NP_002383.2
NCBI GenBank Nucleotide #	NM_002392.5
UniProt Accession #	Q00987; Q13226; Q13297; Q13298; Q13299; Q13300; Q13301; Q53XW0; Q71TW9; Q8WYJ1; A6NL51; A8K2S6
Molecular Weight	55,233 Da
NCBI Official Full Name	E3 ubiquitin-protein ligase Mdm2 isoform a
NCBI Official Synonym Full Names	MDM2 oncogene, E3 ubiquitin protein ligase
NCBI Official Symbol	MDM2
NCBI Official Synonym Symbols	HDMX; hdm2; ACTFS
NCBI Protein Information	E3 ubiquitin-protein ligase Mdm2; oncoprotein Mdm2; Mdm2, p53 E3 ubiquitin protein ligase homolog; double minute 2, human homolog of; p53-binding protein; Mdm2, transformed 3T3 cell double minute 2, p53 binding protein

UniProt Protein Name	E3 ubiquitin-protein ligase Mdm2
UniProt Gene Name	MDM2
UniProt Synonym Gene Names	Hdm2
UniProt Entry Name	MDM2_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/monoclonal/mdm2-recombinant-antibody-clone-14f8-flow-cytometry-immunofluorescence-immunohistochemistry-elisa/279110>