

ERBB2 recombinant antibody

ERBB2 Recombinant Monoclonal Antibody


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA279143	HEK293F Cell	Human, Mouse	Immunohistochemistry, ELISA

PRODUCT SYNONYMS


ERBB2; Monoclonal Recombinant Antibody; ERBB2 Recombinant Monoclonal Antibody; Verb b2 erythroblastic leukemia viral oncogene homolog 2; neuro/glioblastoma derived oncogene homolog; C erb B2/neu protein; CD340; CD340 antigen; Cerb B2/neu protein; CerbB2; Erb b2 receptor tyrosine kinase 2; ErbB-2 proto-oncogene; ERBB2_HUMAN; HER 2; HER 2/NEU; HER2; Herstatin; Human epidermal growth factor receptor 2; Metastatic lymph node gene 19 protein; MLN 19; MLN19; NEU; NEU proto oncogene; Neuro/glioblastoma derived oncogene homolog; Neuroblastoma/glioblastoma derived oncogene homolog; NGL; p185erbB2; Proto-oncogene c-ErbB-2; Proto-oncogene Neu; Receptor tyrosine-protein kinase erbB-2; TKR1; Tyrosine kinase type cell surface receptor HER2; Tyrosine kinase-type cell surface receptor HER2; V erb b2 avian erythroblastic leukemia viral oncogene homolog 2 (neuro/glioblastoma derived oncogene homolog); V erb b2 avian erythroblastic leukemia viral oncogene homolog 2; V erb b2 avian erythroblastic leukemia viral oncoprotein 2; V erb b2 erythroblastic leukemia viral oncogene homolog 2; neuro/glioblastoma derived oncogene homolog (avian); ERBB2 recombinant antibody

PRODUCT SPECIFICATIONS

Host	HEK293F Cell
Reactivity	Human, Mouse
Clonality	Monoclonal
Isotype	Mouse IgG
Clone Number	9E2
Purity/Purification	Affinity purified
Form/Format	Liquid; 0.03% Proclin 300, 50% glycerol, 0.01M PBS, pH 7.4.
Applicable Applications for ERBB2 recombinant antibody	IHC (Immunohistochemistry), ELISA
Immunogen Species	Human
Immunogen	Recombinant Human ERBB2 protein
Conjugation	Non-conjugated
Research Area	Cancer; Tags & Cell Markers; Immunology; Signal transduction
Preparation and Storage	Upon receipt, store at -20 degree C or -80 degree C. Avoid repeated freeze.



IHC (Immunohistochemistry)
IHC image of AAA279143 diluted at 1:600 and staining in paraffin-embedded human testis tissue performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a Goat anti-Mouse IgG labeled by HRP and visualized using 0.05% DAB.



IHC (Immunohistochemistry)
IHC image of AAA279143 diluted at 1:600 and staining in paraffin-embedded human breast cancer tissue performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a Goat anti-Mouse IgG labeled by HRP and visualized using 0.05% DAB.

NCBI AND UNIPROT INFORMATION

NCBI GI #	54792098
NCBI GeneID	2064
NCBI Accession #	NP_001005862.1
NCBI GenBank Nucleotide #	NM_001005862.1
UniProt Accession #	P04626; Q14256; Q6LDV1; Q9UMK4; B2RZG3; B4DHN3
Molecular Weight	137,910 Da
NCBI Official Full Name	receptor tyrosine-protein kinase erbB-2 isoform b
NCBI Official Synonym Full Names	v-erb-b2 erythroblastic leukemia viral oncogene homolog 2, neuro/glioblastoma derived oncogene homolog (avian)
NCBI Official Symbol	ERBB2
NCBI Official Synonym Symbols	NEU; NGL; HER2; TKR1; CD340; HER-2; MLN 19; HER-2/neu
NCBI Protein Information	receptor tyrosine-protein kinase erbB-2; herstatin; p185erbB2; proto-oncogene Neu; c-erb B2/neu protein; proto-oncogene c-ErbB-2; metastatic lymph node gene 19 protein; tyrosine kinase-type cell surface receptor HER2; neuroblastoma/glioblastoma derived on
UniProt Protein Name	Receptor tyrosine-protein kinase erbB-2
UniProt Gene Name	ERBB2
UniProt Synonym Gene Names	HER2; MLN19; NEU; NGL; MLN 19
UniProt Entry Name	ERBB2_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.