

PLAUR recombinant antibody

PLAUR Recombinant Monoclonal Antibody


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA279183	HEK293F Cell	Human	Flow Cytometry, Functional Assay, Immunofluorescence, ELISA

PRODUCT SYNONYMS

PLAUR; Monoclonal Recombinant Antibody; PLAUR Recombinant Monoclonal Antibody; CD 87; CD87; CD87 antigen; MO 3; MO3; Monocyte activation antigen Mo3; Plasminogen activator receptor urokinase; Plasminogen activator urokinase receptor; U PAR; u plasminogen activator receptor; U-PAR; u-plasminogen activator receptor form 2; UPA receptor; uPAR; UPAR_HUMAN; Urinary plasminogen activator receptor; URKR; Urokinase plasminogen activator receptor; Urokinase plasminogen activator surface receptor; urokinase-type plasminogen activator (uPA) receptor; PLAUR recombinant antibody


PRODUCT SPECIFICATIONS

Host	HEK293F Cell
Reactivity	Human
Clonality	Monoclonal
Isotype	Mouse IgG
Clone Number	2E2
Purity/Purification	Affinity purified
Form/Format	Liquid; 0.03% Proclin 300, 50% glycerol, 0.01M PBS, pH 7.4.
Applicable Applications for PLAUR recombinant antibody	FCM/FACS (Flow Cytometry), IF (Immunofluorescence), ELISA
Immunogen Species	Human
Immunogen	Recombinant Human PLAUR protein
Conjugation	Non-conjugated
Research Area	Cancer; Cardiovascular; Immunology
Preparation and Storage	Upon receipt, store at -20 degree C or -80 degree C. Avoid repeated freeze.




FCM/FACS (Flow Cytometry)

Overlay Peak curve showing Hela cells stained with AAA279183 (red line) at 1:100. Then 10% normal goat serum was incubated to block non-specific protein-protein interactions followed by the antibody (1ug/1*10⁶cells) for 45 min at 4°C. The secondary antibody used was FITC-conjugated Goat Anti-Mouse IgG(H+L) at 1/200 dilution for 35 min at 4°C. Isotype control antibody (green line) was mouse IgG1 (1ug/1*10⁶cells) used under the same conditions. Acquisition of >10,026 events was performed.



IF (Immunofluorescence)

Immunofluorescence staining of hela cell with AAA279183 at 1:30, counter-stained with DAPI. The cells were fixed in 4% formaldehyde and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4C. The secondary antibody was FITC-conjugated AffiniPure Goat Anti-Mouse IgG(H+L).



IF (Immunofluorescence)

Immunofluorescence staining of SH-SY5Y cell with AAA279183 at 1:30, counter-stained with DAPI. The cells were fixed in 4% formaldehyde and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4C. The secondary antibody was FITC-conjugated AffiniPure Goat Anti-Mouse IgG(H+L).

NCBI AND UNIPROT INFORMATION

NCBI GI #	4505865
NCBI GeneID	5329
NCBI Accession #	NP_002650.1
NCBI GenBank Nucleotide #	NM_002659.3
UniProt Accession #	Q03405; Q12876; Q15845; Q16887; Q6IB52; Q9BWT0; Q9NYC8; Q9UD69; Q9UEA6; Q9UM92; Q9UMV0; A8K409
Molecular Weight	36,978 Da
NCBI Official Full Name	urokinase plasminogen activator surface receptor isoform 1
NCBI Official Synonym Full Names	plasminogen activator, urokinase receptor
NCBI Official Symbol	PLAUR
NCBI Official Synonym Symbols	CD87; UPAR; URKR; U-PAR
NCBI Protein Information	urokinase plasminogen activator surface receptor; monocyte activation antigen Mo3; u-plasminogen activator receptor form 2; urokinase-type plasminogen activator (uPA) receptor

UniProt Protein Name	Urokinase plasminogen activator surface receptor
UniProt Gene Name	PLAUR
UniProt Synonym Gene Names	MO3; UPAR; U-PAR; uPAR
UniProt Entry Name	UPAR_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/monoclonal/plaur-recombinant-antibody-clone-2e2-flow-cytometry-immunofluorescence-elisa/279183>