

Dickkopf-related protein 1 (DKK1) Active Protein | DKK1 active protein

Recombinant Human Dickkopf-related protein 1 (DKK1)(Active)

CATALOG #
AAA279208

HOST
Mammalian Cell


PRODUCT SYNONYMS

Dickkopf-related protein 1 (DKK1); Recombinant Human Dickkopf-related protein 1 (DKK1)(Active); Dickkopf-1; Dkk-1; hDkk-1; SK; DKK1 active protein


PRODUCT SPECIFICATIONS

Host	Mammalian Cell
Purity/Purification	Greater than 95% as determined by SDS-PAGE.
Form/Format	Lyophilized powder Lyophilized from a 0.2 um filtered 20 mM Tris-HCl, 0.5 M NaCl, 6% Trehalose, pH 8.0
Sequence Positions	32-266aa; Full Length of Mature Protein
Sequence	TLNSVLNSNA IKNLPPPLGG AAGHPGSAVS AAPGILYPGG NKYQTIDNYQ PYPCAEDEEC GTDEYCASPT RGGDAGVQIC LACRKRRKRC MRHAMCCPGN YCKNGICVSS DQNHFRGEIE ETITESFGND HSTLDGYSRR TTLSSKMYHT KGQEGSVCLR SSDCASGLCC ARHFWSKICK PVLKEGQVCT KHRRKGSHGL EIFQRCYCGE GLSCRIQKDH HQASNSSRLH TCQRH\$
Species	Homo sapiens (Human)
Tag	C-terminal 10xHis-tagged
Endotoxin	Less than 1.0 EU/ug as determined by LAL method.
Research Area	Cardiovascular
Biological Activity	Measured by its binding ability in a functional ELISA. Immobilized Human DKK1 at 2 ug/mL can bind Anti-DKK1 recombinant antibody, the EC50 is 1.283-2.544 ng/mL.
Preparation and Storage	The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself. Generally, the shelf life of liquid form is 6 months at -20 degree C/-80 degree C. The shelf life of lyophilized form is 12 months at -20 degree C/-80 degree C. Repeated freezing and thawing is not recommended. Store working aliquots at 4 degree C for up to one week. We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL. We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20 degree C/-80 degree C. Our default final concentration of glycerol is 50%. Customers could use it as reference.

IMAGES



Bioactivity
Measured by its binding ability in a functional ELISA. Immobilized Human DKK1 at 2 ?g/mL can bind Anti-DKK1 recombinant antibody, the EC50 is 1.283-2.544 ng/mL.



SDS-PAGE
Tris-Glycine gel) Discontinuous SDS-PAGE (reduced) with 5% enrichment gel and 15% separation gel.

ADDITIONAL INFORMATION

Related Product Information for DKK1 active protein	Antagonizes canonical Wnt signaling by inhibiting LRP5/6 interaction with Wnt and by forming a ternary complex with the transmembrane protein KREMEN that promotes internalization of LRP5/6.DKKs play an important role in vertebrate development, where they locally inhibit Wnt regulated processes such as antero-posterior axial patterning, limb development, somitogenesis and eye formation. In the adult, Dkks are implicated in bone formation and bone disease, cancer and Alzheimer disease.Inhibits the pro-apoptotic function of KREMEN1 in a Wnt-independent manner, and has anti-apoptotic activity.
Product Categories/Family for DKK1 active protein	Active Protein & In Stock Protein
References	"Regulation of Wnt/LRP signaling by distinct domains of Dickkopf proteins."Brott B.K., Sokol S.Y.Mol Cell Biol 22:6100-6110(2002)

NCBI AND UNIPROT INFORMATION

NCBI GI #	7110719
NCBI GeneID	22943
NCBI Accession #	NP_036374.1
NCBI GenBank Nucleotide #	NM_012242.2
UniProt Accession #	O94907; B2RC19
Molecular Weight	28,672 Da
NCBI Official Full Name	dickkopf-related protein 1
NCBI Official Synonym Full Names	dickkopf WNT signaling pathway inhibitor 1
NCBI Official Symbol	DKK1
NCBI Official Synonym Symbols	SK; DKK-1
NCBI Protein Information	dickkopf-related protein 1; hDkk-1; dickkopf-1 like; dickkopf 1 homolog; dickkopf-like protein 1; dickkopf related protein-1
UniProt Protein Name	Dickkopf-related protein 1
UniProt Gene Name	DKK1
UniProt Synonym Gene Names	Dickkopf-1; Dkk-1; hDkk-1

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/mammalian-cell-active-protein/dickkopf-related-protein-1-dkk1/279208>