

MMAE&MMAF Recombinant Antibody | MMAE recombinant antibody

MMAE&MMAF Recombinant Monoclonal Antibody

CATALOG # AAA279299	HOST HEK293F Cell	REACTIVITY Species Independent	APPLICATIONS ELISA
-------------------------------	-----------------------------	--	------------------------------


PRODUCT SYNONYMS

MMAE&MMAF; Monoclonal Recombinant Antibody; MMAE&MMAF Recombinant Monoclonal Antibody; MMAE; Monomethyl auristatin E; SGD-1010; Vedotin; MMAF; Monomethylauristatin F\$; MMAE recombinant antibody

PRODUCT SPECIFICATIONS


Host	HEK293F Cell
Reactivity	Species Independent
Clonality	Monoclonal
Isotype	IgG2b
Clone Number	25G12B5
Purity/Purification	Affinity-chromatography
Form/Format	Liquid; Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4
Applicable Applications for MMAE recombinant antibody	ELISA
Antigen Species	Human
Antigen	MMAE-BSA
Conjugation	Non-conjugated
Research Area	Others
Preparation and Storage	Upon receipt, store at -20 degree C or -80 degree C. Avoid repeated freeze.

IMAGES




ELISA

The Binding Activity of Disitamab vedotin (Synonyms: RC48) and Depatuxizumab mafodotin (Synonyms: ABT-414) with Anti-MMAE&MMAF antibody. Activity: Measured by its binding ability in a functional ELISA. Immobilized Disitamab vedotin (Synonyms: RC48) and Depatuxizumab mafodotin (Synonyms: ABT-414) at 2 ug/mL can bind Anti-MMAE&MMAF antibody, the EC50 is 3.523 to 5.356 ng/mL and 3.022 to 4.198 ng/mL, respectively.



ELISA

The Binding Activity of Depatuxizumab mafodotin (Synonyms: ABT-414) with Anti-MMAE&MMAF antibody. Activity: Measured by its binding ability in a functional ELISA. Immobilized Depatuxizumab mafodotin (Synonyms: ABT-414) at 2 ug/mL can bind Anti-MMAE&MMAF antibody, the EC50 is 3.022 to 4.198 ng/mL.



ELISA

The Binding Activity of Disitamab vedotin (Synonyms: RC48) with Anti-MMAE&MMAF antibodyActivity: Measured by its binding ability in a functional ELISA. Immobilized Disitamab vedotin (Synonyms: RC48) at 2 ug/mL can bind Anti-MMAE&MMAF antibody, the EC50 is 3.523 to 5.356 ng/mL.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/monoclonal/mmaemmaf-recombinant-antibody-clone-25g12b5-elisa/279299>