

HGF recombinant antibody

HGF Recombinant Monoclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA279317	HEK293F Cell	Human	Immunohistochemistry, ELISA


PRODUCT SYNONYMS

HGF; Monoclonal Recombinant Antibody; HGF Recombinant Monoclonal Antibody; DFN39 antibody; F TCF antibody; Fibroblast derived tumor cytotoxic factor antibody; Hepatocyte growth factor (hepatopoietin A; scatter factor) antibody; Hepatocyte growth factor antibody; Hepatocyte growth factor beta chain antibody; Hepatocyte growth factor precursor antibody; Hepatopoietin A antibody; Hepatopoietin-A antibody; Hgf antibody; HGF_HUMAN antibody; HGFB antibody; HPTA antibody; Lung fibroblast derived mitogen antibody; OTTHUMP00000161349 antibody; OTTHUMP00000206710 antibody; OTTHUMP00000206711 antibody; OTTHUMP00000206712 antibody; OTTHUMP00000206713 antibody; OTTHUMP00000206730 antibody; Scatter factor antibody; SF antibody; HGF recombinant antibody

PRODUCT SPECIFICATIONS


Host	HEK293F Cell
Reactivity	Human
Clonality	Monoclonal
Isotype	IgG1
Clone Number	9H7
Purity/Purification	Affinity-chromatography
Form/Format	Liquid; Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4
Applicable Applications for HGF recombinant antibody	IHC (Immunohistochemistry), ELISA
Antigen Species	Human
Antigen	Recombinant Human HGF protein
Conjugation	Non-conjugated
Research Area	Cancer, Signal transduction
Preparation and Storage	Upon receipt, store at -20 degree C or -80 degree C. Avoid repeated freeze.

IMAGES



IHC (Immunohistochemisry)

IHC image of AAA279317 diluted at 1:100 and staining in paraffin-embedded human kidney tissue performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 3 degree C overnight. The primary is detected by a Goat anti-Human IgG labeled by HRP and visualized using 0.05% DAB.



IHC (Immunohistochemistry)

IHC image of AAA279317 diluted at 1:100 and staining in paraffin-embedded human Liver tissue performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 3 degree C overnight. The primary is detected by a Goat anti-Human IgG labeled by HRP and visualized using 0.05% DAB.

NCBI AND UNIPROT INFORMATION

NCBI GI #	123116
NCBI GeneID	3082
UniProt Accession #	P14210; Q02935; Q13494; Q14519; Q3KRB2; Q8TCE2; Q9BYL9; Q9BYM0; Q9UDU6; A1L3U6
Molecular Weight	728
NCBI Official Full Name	Hepatocyte growth factor
NCBI Official Synonym Full Names	hepatocyte growth factor (hepapoietin A; scatter factor)
NCBI Official Symbol	HGF
NCBI Official Synonym Symbols	SF; HGFB; HPTA; F-TCF; DFNB39
NCBI Protein Information	hepatocyte growth factor; hepatopoeitin-A; hepatopietin-A; lung fibroblast-derived mitogen; fibroblast-derived tumor cytotoxic factor
UniProt Protein Name	Hepatocyte growth factor
UniProt Gene Name	HGF
UniProt Synonym Gene Names	HPTA; SF
UniProt Entry Name	HGF_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.