

AOC3 recombinant antibody

AOC3 Recombinant Monoclonal Antibody

CATALOG # AAA279319	HOST HEK293F Cell	REACTIVITY Human	APPLICATIONS Immunohistochemistry, ELISA
-------------------------------	-----------------------------	----------------------------	--


PRODUCT SYNONYMS

AOC3; Monoclonal Recombinant Antibody; AOC3 Recombinant Monoclonal Antibody; Amine oxidase; copper containing 3 (vascular adhesion protein 1) antibody; AOC3 antibody; AOC3_HUMAN antibody; Copper amine oxidase antibody; HPAO antibody; Membrane copper amine oxidase antibody; Membrane primary amine oxidase antibody; Semicarbazide sensitive amine oxidase antibody; Semicarbazide-sensitive amine oxidase antibody; SSAO antibody; VAP-1 antibody; Vascular adhesion protein 1 antibody; VP97 antibody; AOC3 recombinant antibody


PRODUCT SPECIFICATIONS

Host	HEK293F Cell
Reactivity	Human
Clonality	Monoclonal
Isotype	hIgG4
Clone Number	3D3
Purity/Purification	Affinity-chromatography
Form/Format	Liquid; Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4
Applicable Applications for AOC3 recombinant antibody	IHC (Immunohistochemistry), ELISA
Antigen Species	Human
Antigen	Recombinant Human AOC3 protein
Conjugation	Non-conjugated
Research Area	Neuroscience, Signal transduction
Preparation and Storage	Upon receipt, store at -20 degree C or -80 degree C. Avoid repeated freeze.

IMAGES



IHC (Immunohistochemistry)
IHC image of AAA279319 diluted at 1?150 and staining in paraffin-embedded human adrenal gland tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a Goat anti-human polymer IgG labeled by HRP and visualized using 0.05% DAB.



IHC (Immunohistochemistry)
IHC image of AAA279319 diluted at 1?150 and staining in paraffin-embedded human lung tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a Goat anti-human polymer IgG labeled by HRP and visualized using 0.05% DAB.

NCBI AND UNIPROT INFORMATION

NCBI GI #	2501336
NCBI GeneID	8639
UniProt Accession #	Q16853; Q45F94; B2RCI5; K7ESB3; L0L8N9
Molecular Weight	763
NCBI Official Full Name	Membrane primary amine oxidase
NCBI Official Synonym Full Names	amine oxidase, copper containing 3
NCBI Official Symbol	AOC3
NCBI Official Synonym Symbols	HPAO; SSAO; VAP1; VAP-1
NCBI Protein Information	membrane primary amine oxidase; copper amine oxidase; placenta copper monamine oxidase; semicarbazide-sensitive amine oxidase; amine oxidase, copper containing 3 (vascular adhesion protein 1)
UniProt Protein Name	Membrane primary amine oxidase
UniProt Gene Name	AOC3
UniProt Synonym Gene Names	VAP1; SSAO; VAP-1
UniProt Entry Name	AOC3_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/monoclonal/aoc3-recombinant-antibody-clone-3d3-immunohistochemistry-elisa/279319>