

CSF3R recombinant antibody

CSF3R Recombinant Monoclonal Antibody


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA279324	HEK293F Cell	Human	Immunohistochemistry, ELISA

PRODUCT SYNONYMS


CSF3R; Monoclonal Recombinant Antibody; CSF3R Recombinant Monoclonal Antibody; CD 114 antibody; CD114 antibody; CD114 antigen antibody; Colony stimulating factor 3 receptor (granulocyte) antibody; Colony stimulating factor 3 receptor antibody; CSF 3R antibody; CSF3R antibody; CSF3R_HUMAN antibody; Csfr antibody; G CSF R antibody; G-CSF receptor antibody; G-CSF-R antibody; GCSFR antibody; Granulocyte colony stimulating factor receptor antibody; Granulocyte colony-stimulating factor receptor antibody; OTTHUMP00000009703 antibody; OTTHUMP00000009704 antibody; OTTHUMP00000009705 antibody; CSF3R recombinant antibody

PRODUCT SPECIFICATIONS

Host	HEK293F Cell
Reactivity	Human
Clonality	Monoclonal
Isotype	hIgG1
Clone Number	11C11
Purity/Purification	Affinity-chromatography
Form/Format	Liquid; Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4
Applicable Applications for CSF3R recombinant antibody	IHC (Immunohistochemistry), ELISA
Antigen Species	Human
Antigen	Recombinant Human CSF3R protein
Conjugation	Non-conjugated
Research Area	Cancer, Immunology, Stem cells
Preparation and Storage	Upon receipt, store at -20 degree C or -80 degree C. Avoid repeated freeze.



IHC (Immunohistochemistry)
IHC image of AAA279324 diluted at 1?100 and staining in paraffin-embedded human cervical cancer performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a Goat anti-human polymer IgG labeled by HRP and visualized using 0.05% DAB.



IHC (Immunohistochemistry)
IHC image of AAA279324 diluted at 1?100 and staining in paraffin-embedded human testis tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a Goat anti-human polymer IgG labeled by HRP and visualized using 0.05% DAB.

NCBI AND UNIPROT INFORMATION

NCBI GI #	27437045
NCBI GeneID	1441
NCBI Accession #	NP_758519.1
NCBI GenBank Nucleotide #	NM_172313.2
UniProt Accession #	Q99062
Molecular Weight	92,156 Da
NCBI Official Full Name	granulocyte colony-stimulating factor receptor isoform d
NCBI Official Synonym Full Names	colony stimulating factor 3 receptor (granulocyte)
NCBI Official Symbol	CSF3R
NCBI Official Synonym Symbols	CD114; GCSFR
NCBI Protein Information	granulocyte colony-stimulating factor receptor; G-CSF-R; CD114 antigen; G-CSF receptor
UniProt Protein Name	Granulocyte colony-stimulating factor receptor
UniProt Gene Name	CSF3R
UniProt Synonym Gene Names	GCSFR; G-CSF receptor; G-CSF-R
UniProt Entry Name	CSF3R_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/monoclonal/csf3r-recombinant-antibody-clone-11c11-immunohistochemistry-elisa/279324>