

Interleukin-4 (IL4) Active Protein | IL4 active protein

Recombinant Human Interleukin-4 (IL4)(Active)

CATALOG #
AAA279363

HOST
Mammalian Cell



PRODUCT SYNONYMS

Interleukin-4 (IL4); Recombinant Human Interleukin-4 (IL4)(Active); Interleukin-4; IL-4; B-cell stimulatory factor 1 (BSF-1); Binetrakin; Lymphocyte stimulatory factor 1; Pitrakinra; IL4 active protein

PRODUCT SPECIFICATIONS

Host	Mammalian Cell
Purity/Purification	Greater than 90% as determined by SDS-PAGE.
Form/Format	Lyophilized powder; Lyophilized from a 0.2um sterile filtered PBS, 6% Trehalose, pH 7.4
Sequence	HKCDITLQEI IKTLSLSEQ KTLCTELTVT DIFAASKNTT EKETFCRAAT VLRQFYSHHE KDTRCLGATA QQFHRHKQLI RFLKRLDRNL WGLAGLNSCP VKEANQSTLE NFLERLKTIM REKYSKCSS
Sequence Length	25-153aa; Full Length of Mature Protein
Species	Homo sapiens (Human)
Tag	C-terminal 10xHis-tagged
Endotoxin	Less than 1.0 EU/ug as determined by LAL method.
Biological Activity	The ED50 as determined by the dose-dependent stimulation of the proliferation of TF-1 cells is 1.864-4.949ng/mL.
Reconstitution	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL. We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20 degree C/-80 degree C. Our default final concentration of glycerol is 50%. Customers could use it as reference.
Preparation and Storage	The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself. Generally, the shelf life of liquid form is 6 months at -20 degree C/-80 degree C. The shelf life of lyophilized form is 12 months at -20 degree C/-80 degree C. Notes: Repeated freezing and thawing is not recommended. Store working aliquots at 4 degree C for up to one week.

IMAGES



Activity

SDS-PAGE
Tris-Glycine gel) Discontinuous SDS-PAGE (reduced) with 5% enrichment gel and 15% separation gel.

ADDITIONAL INFORMATION

Related Product Information for IL4 active protein	Cytokine secreted primarily by mast cells, T-cells, eosinophils, and basophils that plays a role in regulating antibody production, hematopoiesis and inflammation, and the development of effector T-cell responses .
Product Categories/Family for IL4 active protein	Immunology
References	An IL-4 signalling axis in bone marrow drives pro-tumorigenic myelopoiesis. LaMarche N.M., Hegde S., Park M.D., Maier B.B., Troncoso L., Le Berichel J., Hamon P., Belabel M., Mattiuz R., Merad M.Nature 625:166-174 (2024)

NCBI AND UNIPROT INFORMATION

NCBI GI #	4504669
NCBI GeneID	3565
NCBI Accession #	NP_000580.1
NCBI GenBank Nucleotide #	NM_000589.3
UniProt Accession #	P05112; Q14630
Molecular Weight	17,492 Da
NCBI Official Full Name	interleukin-4 isoform 1
NCBI Official Synonym Full Names	interleukin 4
NCBI Official Symbol	IL4
NCBI Official Synonym Symbols	BSF1; IL-4; BCGF1; BSF-1; BCGF-1
NCBI Protein Information	interleukin-4; binetrakin; pitrakinra; B cell growth factor 1; B_cell stimulatory factor 1; lymphocyte stimulatory factor 1
UniProt Protein Name	Interleukin-4
UniProt Gene Name	IL4
UniProt Synonym Gene Names	IL-4; BSF-1
UniProt Entry Name	IL4_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

