

Interleukin-15 receptor subunit alpha (IL15RA) Active Protein | IL15RA active protein

Recombinant Human Interleukin-15 receptor subunit alpha (IL15RA), partial (Active)

CATALOG #
AAA279381


HOST
Mammalian Cell

PRODUCT SYNONYMS

Interleukin-15 receptor subunit alpha (IL15RA); Recombinant Human Interleukin-15 receptor subunit alpha (IL15RA); partial (Active); IL-15 receptor subunit alpha; IL-15R-alpha; IL-15RA; IL15RA active protein

PRODUCT SPECIFICATIONS

Host	Mammalian Cell
Purity/Purification	Greater than 95% as determined by SDS-PAGE.
Form/Format	Lyophilized powder; Lyophilized from a 0.2um sterile filtered PBS, 6% Trehalose, pH 7.4
Sequence	ITCPPMSVE HADIWVKSYS LYSRERYICN SGFKRKAGTS SLTECVLNKA TNVAHWTTPS LKCIRDPALV HQRPAAPPSTV TTAGVTPQPE SLSPSGKEPA ASSPSSNNTA ATTAIVPGS QLMPKSPST GTTEISSHES SHGTPSQTTA KNWELTASAS HQPPGVYPQG HSDTTVAIST STVLLCGLSA VSLLACYLKS RQTPPLASVE MEAMEALPVT WGTSS
Sequence Length	31-205aa; Partial
Species	Homo sapiens (Human)
Tag	C-terminal 10xHis-tagged
Endotoxin	Less than 1.0 EU/ug as determined by LAL method.
Biological Activity	Measured by its binding ability in a functional ELISA. Immobilized Human IL15RA at 2ug/mL can bind Human IL15. The EC50 is 32.53-38.41ng/mL.
Reconstitution	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL. We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20 degree C/-80 degree C. Our default final concentration of glycerol is 50%. Customers could use it as reference.
Preparation and Storage	The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself. Generally, the shelf life of liquid form is 6 months at -20 degree C/-80 degree C. The shelf life of lyophilized form is 12 months at -20 degree C/-80 degree C. Notes: Repeated freezing and thawing is not recommended. Store working aliquots at 4 degree C for up to one week.



SDS-PAGE
Tris-Glycine gel) Discontinuous SDS-PAGE (reduced) with 5% enrichment gel and 15% separation gel.

ADDITIONAL INFORMATION

Related Product Information for IL15RA active protein	High-affinity receptor for interleukin-15.Can signal both in cis and trans where IL15R from one subset of cells presents IL15 to neighboring IL2RG-expressing cells.In neutrophils, binds and activates kinase SYK in response to IL15 stimulation.In neutrophils, required for IL15-induced phagocytosis in a SYK-dependent manner.Expression of different isoforms may alter or interfere with signal transduction.
References	Complete sequencing and characterization of 21,243 full-length human cDNAs. Ota T., Suzuki Y., Nishikawa T., Otsuki T., Sugiyama T., Irie R., Wakamatsu A., Hayashi K., Sato H., Sugano S.Nat. Genet. 36:40-45 (2004)

NCBI AND UNIPROT INFORMATION

NCBI GI #	343780891
NCBI GeneID	3601
NCBI Accession #	NP_001230468.1
NCBI GenBank Nucleotide #	NM_001243539.1
UniProt Accession #	Q13261; Q5JVA1; Q5JVA2; Q5JVA4; Q6B0J2; Q7LDR4; Q7Z609; B4E2C2
Molecular Weight	28,233 Da
NCBI Official Full Name	interleukin-15 receptor subunit alpha isoform 3
NCBI Official Synonym Full Names	interleukin 15 receptor, alpha
NCBI Official Symbol	IL15RA
NCBI Official Synonym Symbols	CD215
NCBI Protein Information	interleukin-15 receptor subunit alpha
UniProt Protein Name	Interleukin-15 receptor subunit alpha
UniProt Gene Name	IL15RA
UniProt Synonym Gene Names	IL-15 receptor subunit alpha; IL-15R-alpha; IL-15RA; sIL-15 receptor subunit alpha; sIL-15R-alpha; sIL-15RA
UniProt Entry Name	I15RA_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.