

C-C chemokine receptor type 5 (CCR5)-VLPs Recombinant Protein | CCR5-VLPs recombinant protein

Recombinant Human C-C chemokine receptor type 5 (CCR5)-VLPs (Active)

CATALOG # AAA279393	HOST Mammalian cell
-------------------------------	-------------------------------


PRODUCT SYNONYMS

C-C chemokine receptor type 5 (CCR5)-VLPs; Recombinant Human C-C chemokine receptor type 5 (CCR5)-VLPs (Active); C-C CKR-5; CC-CKR-5; CCR-5; CCR5; CHEMR13; HIV-1 fusion coreceptor; CD antigen CD195; CCR5-VLPs recombinant protein

PRODUCT SPECIFICATIONS

Host	Mammalian cell
Purity/Purification	The purity information is not available for VLPs proteins.
Form/Format	Lyophilized powder. Lyophilized from a 0.2um sterile filtered PBS, 6% Trehalose, pH 7.4.
Sequence	MDYQVSSPIY DINYTSEPC QKINVKQIAA RLLPPLYSLV FIFGFVGNML VILILINCKR LKSMTDIYLL NLAISDLFFL LTVPFWAHYA AAQWDFGNTM CQLLTGLYFI GFFSGIFFII LLTIDRYLAV VHAVFALKAR TVTFGVVTSTV ITWVVAVFAS LPGIIFTRSQ KEGLHYTCSS HFPYSQYQFW KNFQTLKIVI LGLVLP LLVM VICYSGLKLT LLRCRNEKKR HRAVRLIFTI MIVYFLFWAP YNIVLLLNTF QEFFGLNNCS SSNRLDQAMQ VTETLGMTHC CINPIIYAFV GEKFRNYLLV FFQKHIKRF CKCCSIFQQE APERASSVYT RSTGEQEISV GL
Sequence Length	1-352aa; Full Length
Species	Homo sapiens (Human)
Tag	C-terminal 10xHis-tagged (This tag can be tested only under denaturing conditions)
Transmembrane Domain	7TM
Endotoxin	Less than 1.0 EU/ug as determined by LAL method.
Reconstitution	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL. Aliquot for long-term storage at -80 degree C. Solubilize for 60 minutes at room temperature with occasional gentle mixing. Avoid vigorous shaking or vortexing.
Biological Activity	Measured by its binding ability in a functional ELISA. Immobilized Human CCR5 at 10ug/mL can bind Anti-CCR5 recombinant antibody . The EC50 is 1.099-1.287ng/mL. The VLPs is negative control.
Preparation and Storage	The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself. Generally, the shelf life of liquid form is 6 months at -20 degree C/-80 degree C. The shelf life of lyophilized form is 12 months at -20 degree C/-80 degree C. Notes: The VLPs are expressed from human 293 cells (HEK293). Mix the sample gently by repeatedly pipetting it up and down. Do not vortex. Repeated freezing and thawing is not recommended. Store the protein at -20 degree C/-80 degree C upon receiving it, and ensure to avoid repeated freezing and thawing, otherwise, it will affect the protein activity. The immunization strategy should be optimized (antigen dose, regimen and adjuvant).

IMAGES



Application Data
Detected by Mouse anti-6*His monoclonal antibody.(This tag can be tested only under denaturing conditions.

ADDITIONAL INFORMATION

Related Product Information for CCR5-VLPs recombinant protein	Receptor for a number of inflammatory CC-chemokines including CCL3/MIP-1-alpha, CCL4/MIP-1-beta and RANTES and subsequently transduces a signal by increasing the intracellular calcium ion level. May play a role in the control of granulocytic lineage proliferation or differentiation. Participates in T-lymphocyte migration to the infection site by acting as a chemotactic receptor. (Microbial infection) Acts as a coreceptor (CD4 being the primary receptor) of human immunodeficiency virus-1/HIV-1.
Product Categories/Family for CCR5-VLPs recombinant protein	Cancer
References	The DNA sequence, annotation and analysis of human chromosome 3.Muzny D.M., Scherer S.E., Kaul R., Wang J., Yu J., Sudbrak R., Buhay C.J., Chen R., Cree A.Nature 440:1194-1198 (2006)

NCBI AND UNIPROT INFORMATION

NCBI GI #	4502639
NCBI GeneID	1234
NCBI Accession #	NP_000570.1
NCBI GenBank Nucleotide #	NM_000579.3
UniProt Accession #	P51681; O14692; O14693; O14695; O14696; O14697; O14698; O14699; O14700; O14701; O14702; O14703
Molecular Weight	40,524 Da
NCBI Official Full Name	C-C chemokine receptor type 5
NCBI Official Synonym Full Names	chemokine (C-C motif) receptor 5 (gene/pseudogene)
NCBI Official Symbol	CCR5
NCBI Official Synonym Symbols	CKR5; CCR-5; CD195; CKR-5; CCCR5; CMKBR5; IDDM22; CC-CKR-5
NCBI Protein Information	C-C chemokine receptor type 5; chemr13; HIV-1 fusion coreceptor; chemokine receptor CCR5; C-C motif chemokine receptor 5 A159A
UniProt Protein Name	C-C chemokine receptor type 5
UniProt Gene Name	CCR5
UniProt Synonym Gene Names	CMKBR5; C-C CKR-5; CC-CKR-5; CCR-5; CCR5
UniProt Entry Name	CCR5_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/ccr5-recombinant-protein-mammalian-cell/279393>