

Rabbit anti-Human, Mouse SMAD4 Polyclonal Antibody | anti-SMAD4 antibody

SMAD4 Polyclonal Antibody

CATALOG # AAA279447	HOST Rabbit	REACTIVITY Human, Mouse	APPLICATIONS Immunofluorescence, Immunohistochemistry, Western Blot
-------------------------------	-----------------------	-----------------------------------	---

PRODUCT SYNONYMS

SMAD4; Polyclonal Antibody; SMAD4 Polyclonal Antibody; JIP; DPC4; MADH4; MYHRS; anti-SMAD4 antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Affinity Purification
Form/Format	PBS with 0.05% proclin300,50% glycerol,pH7.3.
Concentration	1mg/ml (varies by lot)
Sequence Length	552
Applicable Applications for anti-SMAD4 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Species	Human
Immunogen	Recombinant Protein
Immunogen	Recombinant protein of human SMAD4
Calculated Molecular Weight	60kDa
Preparation and Storage	Store at -20 degree C (regular) or -80 degree C (long term). Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

IMAGES



IF (Immunofluorescence)

Immunofluorescence analysis of U2OS cells using SMAD4 antibody. Blue: DAPI for nuclear staining.



WB (Western Blot)

Western blot analysis of extracts of various cell lines, using SMAD4 antibody at 1:1000 dilution.Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution.Lysates/proteins: 25ug per lane.Blocking buffer: 3% nonfat dry milk in TBST.Detection: ECL Basic Kit.Exposure time: 30s.

ADDITIONAL INFORMATION

Related Product Information for anti-SMAD4 antibody	This gene encodes a member of the Smad family of signal transduction proteins. Smad proteins are phosphorylated and activated by transmembrane serine-threonine receptor kinases in response to TGF-beta signaling. The product of this gene forms homomeric complexes and heteromeric complexes with other activated Smad proteins, which then accumulate in the nucleus and regulate the transcription of target genes. This protein binds to DNA and recognizes an 8-bp palindromic sequence (GTCTAGAC) called the Smad-binding element (SBE). The Smad proteins are subject to complex regulation by post-translational modifications. Mutations or deletions in this gene have been shown to result in pancreatic cancer, juvenile polyposis syndrome, and hereditary hemorrhagic telangiectasia syndrome.
Product Categories/Family for anti-SMAD4 antibody	Polyclonal

NCBI AND UNIPROT INFORMATION

NCBI GI #	13959561
NCBI GeneID	4089
UniProt Accession #	Q13485; A8K405
Molecular Weight	552
NCBI Official Full Name	Mothers against decapentaplegic homolog 4
NCBI Official Synonym Full Names	SMAD family member 4
NCBI Official Symbol	SMAD4
NCBI Official Synonym Symbols	JIP; DPC4; MADH4; MYHRS
NCBI Protein Information	mothers against decapentaplegic homolog 4; MAD homolog 4; SMAD, mothers against DPP homolog 4; deleted in pancreatic carcinoma locus 4; deletion target in pancreatic carcinoma 4; mothers against decapentaplegic, Drosophila, homolog of, 4
UniProt Protein Name	Mothers against decapentaplegic homolog 4
UniProt Gene Name	SMAD4
UniProt Synonym Gene Names	DPC4; MADH4; MAD homolog 4; Mothers against DPP homolog 4; SMAD 4; Smad4; hSMAD4
UniProt Entry Name	SMAD4_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/smad4-antibody-immunofluorescence-immunohistochemistry-western-blot-elisa/279447>