

Rabbit DARS Polyclonal Antibody | anti-DARS antibody

DARS Polyclonal Antibody

CATALOG # AAA279770	HOST Rabbit	REACTIVITY Human, Mouse, Rat	APPLICATIONS Immunohistochemistry, Western Blot
-------------------------------	-----------------------	--	---


PRODUCT SYNONYMS

DARS; Polyclonal Antibody; DARS Polyclonal Antibody; HBSL; aspRS; anti-DARS antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Affinity Purification
Concentration	1mg/ml (varies by lot)
Sequence Length	501
Applicable Applications for anti-DARS antibody	IHC (Immunohistochemistry), WB (Western Blot)
Species	Human
Immunogen	Recombinant Protein
Immunogen	Recombinant protein of human DARS
Calculated Molecular Weight	57kDa
Preparation and Storage	Store at -20 degree C (regular) or -80 degree C (long term). Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

IMAGES




IP (Immunoprecipitation)

Immunoprecipitation analysis of 200ug extracts of 293T cells using 1ug DARS antibody. Western blot was performed from the immunoprecipitate using DARS antibody at a dilution of 1:1000.




IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded mouse brain using DARS antibody at dilution of 1:100 (40x lens).




IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded mouse spinal cord using DARS antibody at dilution of 1:100 (40x lens).



IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded rat brain using DARS antibody at dilution of 1:100 (40x lens).



WB (Western Blot)

Western blot analysis of extracts of various cell lines, using DARS Antibody at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 90s.

ADDITIONAL INFORMATION

Related Product Information for anti-DARS antibody	This gene encodes a member of a multienzyme complex that functions in mediating the attachment of amino acids to their cognate tRNAs. The encoded protein ligates L-aspartate to tRNA(Asp). Mutations in this gene have been found in patients showing hypomyelination with brainstem and spinal cord involvement and leg spasticity. Alternative splicing results in multiple transcript variants.
Product Categories/Family for anti-DARS antibody	Polyclonal

NCBI AND UNIPROT INFORMATION

NCBI GI #	20178330
NCBI GeneID	1615
UniProt Accession #	P14868; Q2TNI3; Q32Q69; Q53HV4; Q53YC5; Q68CR9; Q9BW52; A8K3J2; D3DP77
Molecular Weight	501
NCBI Official Full Name	Aspartate--tRNA ligase, cytoplasmic
NCBI Official Synonym Full Names	aspartyl-tRNA synthetase
NCBI Official Symbol	DARS
NCBI Official Synonym Symbols	HBSL; aspRS

NCBI Protein Information	aspartate--tRNA ligase, cytoplasmic; aspartate tRNA ligase 1, cytoplasmic; aspartyl-tRNA synthetase, cytoplasmic; cell proliferation-inducing gene 40 protein
UniProt Protein Name	Aspartate--tRNA ligase, cytoplasmic
UniProt Gene Name	DARS
UniProt Synonym Gene Names	AspRS
UniProt Entry Name	SYDC_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/dars-antibody-immunohistochemistry-western-blot-elisa/279770>