

Rabbit FMOD Polyclonal Antibody | anti-FMOD antibody

FMOD Polyclonal Antibody

CATALOG # AAA279944	HOST Rabbit	REACTIVITY Human, Mouse, Rat	APPLICATIONS Immunohistochemistry, Western Blot
-------------------------------	-----------------------	--	---


PRODUCT SYNONYMS

FMOD; Polyclonal Antibody; FMOD Polyclonal Antibody; FM; SLRR2E; anti-FMOD antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Affinity Purification
Concentration	1mg/ml (varies by lot)
Sequence Length	376
Applicable Applications for anti-FMOD antibody	IHC (Immunohistochemistry), WB (Western Blot)
Species	Human
Immunogen	Recombinant Protein
Immunogen	Recombinant protein of human FMOD
Calculated Molecular Weight	43kDa
Preparation and Storage	Store at -20 degree C (regular) or -80 degree C (long term). Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

IMAGES




IHC (Immunohistochemisry)

Immunohistochemistry of paraffin-embedded mouse testis using FMOD antibody at dilution of 1:100 (20x lens).



IHC (Immunohiostchemistry)

Immunohistochemistry of paraffin-embedded rat testis using FMOD antibody at dilution of 1:100 (20x lens).



WB (Western Blot)

Western blot analysis of extracts of various cell lines, using FMOD antibody at 1:1000 dilution.Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution.Lysates/proteins: 25ug per lane.Blocking buffer: 3% nonfat dry milk in TBST.Detection: ECL Basic Kit.Exposure time: 90s.

ADDITIONAL INFORMATION

Related Product Information for anti-FMOD antibody	Fibromodulin belongs to the family of small interstitial proteoglycans. The encoded protein possesses a central region containing leucine-rich repeats with 4 keratan sulfate chains, flanked by terminal domains containing disulphide bonds. Owing to the interaction with type I and type II collagen fibrils and in vitro inhibition of fibrillogenesis, the encoded protein may play a role in the assembly of extracellular matrix. It may also regulate TGF-beta activities by sequestering TGF-beta into the extracellular matrix. Sequence variations in this gene may be associated with the pathogenesis of high myopia. Alternative splicing results in multiple transcript variants.
Product Categories/Family for anti-FMOD antibody	Polyclonal

NCBI AND UNIPROT INFORMATION

NCBI GI #	223590208
NCBI GeneID	2331
UniProt Accession #	Q06828; Q15331; Q8IV47
Molecular Weight	376
NCBI Official Full Name	Fibromodulin
NCBI Official Synonym Full Names	fibromodulin
NCBI Official Symbol	FMOD
NCBI Official Synonym Symbols	FM; SLRR2E
NCBI Protein Information	fibromodulin; KSPG fibromodulin; collagen-binding 59 kDa protein; keratan sulfate proteoglycan fibromodulin
UniProt Protein Name	Fibromodulin
UniProt Gene Name	FMOD
UniProt Synonym Gene Names	FM; SLRR2E; FM; KSPG fibromodulin
UniProt Entry Name	FMOD_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/fmod-antibody-immunohistochemistry-western-blot-elisa/279944>