

Rabbit SRSF9 Polyclonal Antibody | anti-SRSF9 antibody

SRSF9 Polyclonal Antibody

CATALOG # AAA280102	HOST Rabbit	REACTIVITY Human, Mouse, Rat, Hamster	APPLICATIONS Immunoprecipitation, Western Blot
-------------------------------	-----------------------	---	--

PRODUCT SYNONYMS

SRSF9; Polyclonal Antibody; SRSF9 Polyclonal Antibody; SFRS9; SRp30c; anti-SRSF9 antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat, Hamster
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Affinity Purification
Concentration	1mg/ml (varies by lot)
Sequence Length	221
Applicable Applications for anti-SRSF9 antibody	IP (Immunoprecipitation), WB (Western Blot)
Species	Human
Immunogen	Recombinant Protein
Immunogen	Recombinant protein of human SRSF9
Calculated Molecular Weight	25kDa
Preparation and Storage	Store at -20 degree C (regular) or -80 degree C (long term). Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

IMAGES




IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded mouse heart using SRSF9 antibody at dilution of 1:100 (40x lens).




IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded human stomach using SRSF9 antibody at dilution of 1:100 (40x lens).



IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded human stomach using SRSF9 antibody (40x lens).



IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded mouse heart using SRSF9 antibody (40x lens).



WB (Western Blot)

Western blot analysis of extracts of various cell lines, using SRSF9 antibody at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 10s.

ADDITIONAL INFORMATION

Related Product Information for anti-SRSF9 antibody	The protein encoded by this gene is a member of the serine/arginine (SR)-rich family of pre-mRNA splicing factors, which constitute part of the spliceosome. Each of these factors contains an RNA recognition motif (RRM) for binding RNA and an RS domain for binding other proteins. The RS domain is rich in serine and arginine residues and facilitates interaction between different SR splicing factors. In addition to being critical for mRNA splicing, the SR proteins have also been shown to be involved in mRNA export from the nucleus and in translation. Two pseudogenes, one on chromosome 15 and the other on chromosome 21, have been found for this gene.
Product Categories/Family for anti-SRSF9 antibody	Polyclonal

NCBI AND UNIPROT INFORMATION

NCBI GI #	3929377
NCBI GeneID	8683
UniProt Accession #	Q13242; Q52LD1
Molecular Weight	221
NCBI Official Full Name	Serine/arginine-rich splicing factor 9
NCBI Official Synonym Full Names	serine/arginine-rich splicing factor 9
NCBI Official Symbol	SRSF9
NCBI Official Synonym Symbols	SFRS9; SRp30c
NCBI Protein Information	serine/arginine-rich splicing factor 9; SR splicing factor 9; pre-mRNA-splicing factor SRp30C; splicing factor, arginine/serine-rich 9
UniProt Protein Name	Serine/arginine-rich splicing factor 9
UniProt Gene Name	SRSF9
UniProt Synonym Gene Names	SFRS9; SRP30C
UniProt Entry Name	SRSF9_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/srsf9-antibody-immunoprecipitation-western-blot-elisa/280102>