

Rabbit ACY1 Polyclonal Antibody | anti-ACY1 antibody

ACY1 Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA280122	Rabbit	Human, Mouse, Rat	Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

ACY1; Polyclonal Antibody; ACY1 Polyclonal Antibody; ACY-1; ACY1D; HEL-S-5; anti-ACY1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Affinity Purification
Concentration	1mg/ml (varies by lot)
Sequence Length	408
Applicable Applications for anti-ACY1 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Species	Human
Immunogen	Recombinant Protein
Immunogen	Recombinant protein of human ACY1
Calculated Molecular Weight	45kDa
Preparation and Storage	Store at -20 degree C (regular) or -80 degree C (long term). Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

IMAGES



IF (Immunofluorescence)

Immunofluorescence analysis of HeLa cells using ACY1 antibody. Blue: DAPI for nuclear staining.



WB (Western Blot)

Western blot analysis of extracts of various cell lines, using ACY1 antibody at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 90s.

ADDITIONAL INFORMATION

Related Product Information for anti-ACY1 antibody	This gene encodes a cytosolic, homodimeric, zinc-binding enzyme that catalyzes the hydrolysis of acylated L-amino acids to L-amino acids and an acyl group, and has been postulated to function in the catabolism and salvage of acylated amino acids. This gene is located on chromosome 3p21.1, a region reduced to homozygosity in small-cell lung cancer (SCLC), and its expression has been reported to be reduced or undetectable in SCLC cell lines and tumors. The amino acid sequence of human aminoacylase-1 is highly homologous to the porcine counterpart, and this enzyme is the first member of a new family of zinc-binding enzymes. Mutations in this gene cause aminoacylase-1 deficiency, a metabolic disorder characterized by central nervous system defects and increased urinary excretion of N-acetylated amino acids. Alternative splicing of this gene results in multiple transcript variants. Read-through transcription also exists between this gene and the upstream ABHD14A (abhydrolase domain containing 14A) gene, as represented in GeneID:100526760. A related pseudogene has been identified on chromosome 18.
Product Categories/Family for anti-ACY1 antibody	Polyclonal

NCBI AND UNIPROT INFORMATION

NCBI GI #	461466
NCBI GeneID	95
UniProt Accession #	Q03154; C9J6I6; C9J9D8; C9JWD4
Molecular Weight	408
NCBI Official Full Name	Aminoacylase-1
NCBI Official Synonym Full Names	aminoacylase 1
NCBI Official Symbol	ACY1
NCBI Official Synonym Symbols	ACY-1; ACY1D; HEL-S-5
NCBI Protein Information	aminoacylase-1; epididymis secretory protein Li 5; N-acyl-L-amino-acid amidohydrolase
UniProt Protein Name	Aminoacylase-1
UniProt Gene Name	ACY1
UniProt Synonym Gene Names	ACY-1
UniProt Entry Name	ACY1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/acy1-antibody-immunofluorescence-immunohistochemistry-western-blot-elisa/280122>