

Rabbit PSMD13 Polyclonal Antibody | anti-PSMD13 antibody

PSMD13 Polyclonal Antibody

CATALOG # AAA280236	HOST Rabbit	REACTIVITY Human, Mouse, Rat	APPLICATIONS Immunofluorescence, Immunohistochemistry, Western Blot
-------------------------------	-----------------------	--	---


PRODUCT SYNONYMS

PSMD13; Polyclonal Antibody; PSMD13 Polyclonal Antibody; S11; Rpn9; p40.5; HSPC027; anti-PSMD13 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Affinity Purification
Concentration	1mg/ml (varies by lot)
Sequence Length	376
Applicable Applications for anti-PSMD13 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Species	Human
Immunogen	Recombinant Protein
Immunogen	Recombinant protein of human PSMD13
Calculated Molecular Weight	43kDa
Preparation and Storage	Store at -20 degree C (regular) or -80 degree C (long term). Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.


IMAGES


IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded mouse lung using PSMD13 antibody at dilution of 1:100 (40x lens).


IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded human liver injury using PSMD13 antibody at dilution of 1:100 (40x lens).



WB (Western Blot)

Western blot analysis of extracts of various cell lines, using PSMD13 antibody at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 90s.

ADDITIONAL INFORMATION

Related Product Information for anti-PSMD13 antibody

The 26S proteasome is a multicatalytic proteinase complex with a highly ordered structure composed of 2 complexes, a 20S core and a 19S regulator. The 20S core is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. The 19S regulator is composed of a base, which contains 6 ATPase subunits and 2 non-ATPase subunits, and a lid, which contains up to 10 non-ATPase subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. This gene encodes a non-ATPase subunit of the 19S regulator. Two transcripts encoding different isoforms have been described.

Product Categories/Family for anti-PSMD13 antibody

Polyclonal

NCBI AND UNIPROT INFORMATION

NCBI GI #	317373273
NCBI GeneID	5719
UniProt Accession #	Q9UNM6; O75831; Q53XU2; Q9UNV3; B3KT15
Molecular Weight	376
NCBI Official Full Name	26S proteasome non-ATPase regulatory subunit 13
NCBI Official Synonym Full Names	proteasome (prosome, macropain) 26S subunit, non-ATPase, 13
NCBI Official Symbol	PSMD13
NCBI Official Synonym Symbols	S11; Rpn9; p40.5; HSPC027
NCBI Protein Information	26S proteasome non-ATPase regulatory subunit 13; 26S proteasome subunit p40.5; 26S proteasome regulatory subunit S11; 26S proteasome regulatory subunit RPN9; 26S proteasome regulatory subunit p40.5
UniProt Protein Name	26S proteasome non-ATPase regulatory subunit 13
UniProt Gene Name	PSMD13
UniProt Entry Name	PSD13_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/psmd13-antibody-immunofluorescence-immunohistochemistry-western-blot-elisa/280236>